

# Convincing evidence supports reducing saturated fat to decrease cardiovascular disease risk

Penny M Kris-Etherton,<sup>1</sup> Kristina Petersen,<sup>1</sup> Linda Van Horn<sup>2</sup>

<sup>1</sup>Department of Nutritional Sciences, The Pennsylvania State University, University Park, Pennsylvania, USA

<sup>2</sup>Feinberg School of Medicine, Northwestern University Medical School, Chicago, Illinois, USA

## Correspondence to

Dr Penny M Kris-Etherton; pmk3@psu.edu

Received 7 August 2018

Revised 11 October 2018

Accepted 12 October 2018

## INTRODUCTION

Treatment guidelines and population-based recommendations evolve from research vetted by the scientific community. Healthcare providers require practice guidelines resulting from rigorous review of the totality of evidence. Open discussion/debate among experts is fundamental and encouraged, but blatant disregard for scientific process can lead to confusion and public distrust. A recent example is the controversy among scientists, healthcare professionals and the public about dietary saturated fat (SFA) recommendations to lower risk of cardiovascular disease (CVD).

## Dietary recommendations for SFA from authoritative organisations

More than 60 years ago (in 1957), the American Heart Association (AHA) proposed that modifying dietary fat could reduce the incidence of coronary heart disease (CHD). This was based on research, including controlled feeding studies, and endpoint measures of blood lipids and blood pressure.<sup>1</sup> In 1980, the first Dietary Guidelines for Americans recommended 'avoiding too much SFA'.<sup>2</sup> For decades, research has substantiated this consistent dietary recommendation from authoritative organisations to decrease SFA to reduce CVD risk.<sup>3–6</sup> Current dietary guidelines for SFA and evidence ratings are summarised in table 1. While the strong evidence to reduce SFA specifically targets decreasing low-density lipoprotein (LDL)-cholesterol, it is very much germane to CVD risk reduction.

## Challenging practice guidelines and population-based recommendations: the SFA SAGA

In 2010, a meta-analysis of 21 prospective epidemiological studies totalling 347 747 subjects reported that higher intake of SFA was not associated with an increased risk of CHD, stroke or CVD.<sup>7</sup> Aspects of this methodology were widely criticised,<sup>8–10</sup> most notably the failure to account for the proportional

differences in dietary macronutrient composition with varying SFA intake.<sup>8 10</sup> Using similar meta-analytic methods, and again without considering the macronutrient composition of diets differing in SFA, Chowdhury *et al* and also de Souza *et al* further concluded that SFA was not associated with CHD or CVD or CHD mortality.<sup>11 12</sup> Additional analyses of prospective cohort studies and randomised controlled trials that carefully evaluated both the dietary nutrient profile and SFA intake have consistently demonstrated a lower risk of CHD when polyunsaturated fatty acids (PUFAs) are consumed instead of SFA.<sup>13 14</sup> More recently,<sup>15</sup> findings from the Nurses' Health Study (NHS) (1980–2010) and Health Professionals Follow-up Study (1986–2010) suggested that lowering energy from SFA by 5% and proportionally increasing PUFA, monounsaturated fatty acids (MUFAs) or carbohydrates from whole grains was associated with a 25%, 15% and 9% lower risk of CHD, respectively (PUFAs: HR: 0.75, 95% CI 0.67 to 0.84; p<0.0001; MUFAs: HR: 0.85, 95% CI 0.74 to 0.97; p=0.02; carbohydrates from whole grains: HR: 0.91, 95% CI 0.85 to 0.98; p=0.01).<sup>15</sup> Similarly, lower intake of predominant long-chain SFA in favour of other macronutrients and plant proteins reduced risk of CHD in the NHS and the Health Professionals Follow-up Study.<sup>16</sup> Consumption of 1% of energy from plant proteins instead of C12:0–C18:0 SFA decreased CHD risk (HR: 0.93, 95% CI 0.89 to 0.97; p=0.001).

The clinical trial evidence has focused on evaluating replacement of SFA with PUFA. Based on three recent meta-analyses of clinical studies,<sup>17–19</sup> the debate continues. The first two papers concluded that the evidence does not support current recommendations to replace SFA with PUFA since there was no effect on CHD mortality and total/all-cause mortality.<sup>17 18</sup> Hamley<sup>18</sup> also reported no effect on major CHD events and total CHD



© Author(s) (or their employer(s)) 2018. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Kris-Etherton PM, Petersen K, Van Horn L. *bmjnp* Epub ahead of print: [please include Day Month Year]. doi:10.1136/bmjnp-2018-000009

**Table 1** Authoritative recommendations for saturated fat and evidence ratings

Recommendation			Evidence rating	
2013 AHA/ACC guideline on lifestyle management to reduce cardiovascular risk				
	NHLBI grade	NHLBI evidence statements	ACC/AHA COR	ACC/AHA COR
Aim for a dietary pattern that achieves 5%–6% of calories from SFA	A (strong)*	Evidence statement 11 – strength of evidence: high†	I‡	A§
Reduce per cent of calories from SFA	A (strong)*	Evidence statement 11 – strength of evidence: high†	I‡	A§
2015 National Lipid Association recommendations for patient-centred management of dyslipidaemia: part 2				
	Grade	Strength of recommendation		
Dietary SFA may be partially replaced with unsaturated fats (MUFA and PUFA), as well as proteins, to reach a goal of <7% of energy from SFA.	A*	Moderate¶		
Scientific report of the 2015 Dietary Guidelines Advisory Committee			DGAC grade	
Strong and consistent evidence from RCTs and statistical modelling in prospective cohort studies shows that replacing SFA with PUFA reduces the risk of CVD events and coronary mortality.			Strong**	
2016 ESC/EAS guidelines for the management of dyslipidaemias				
	Magnitude of the effect		Level of evidence	
For SFA, consumption should be <10% of the total caloric intake and should be further reduced (<7% of energy) in the presence of hypercholesterolaemia.	+++††		A‡‡	

Strength of recommendation – strong: there is high certainty, based on evidence, that the net benefit is substantial.

Evidence Statement 11: when food was supplied to adults in a dietary pattern that achieved a macronutrient composition of 5%–6% saturated fat, 26%–27% total fat, 15%–18% protein and 55%–59% carbohydrate compared with the control diet (14%–15% saturated fat, 34%–38% total fat, 13%–15% protein and 48%–51% carbohydrate), LDL-C was lowered 11–13 mg/dL in two studies and 11% in another study. Strength of evidence: high.

Classification of recommendation (COR): class I: benefit >>> risk; procedure/treatment should be performed/administered.

Level of evidence: level A: multiple populations evaluated; data derived from multiple randomised clinical trials or meta-analyses.

RCTs with minor limitations affecting confidence in, or applicability of, the results. Well-designed, well-executed non-randomised controlled studies and well-designed, well-executed observational studies. Well-conducted meta-analyses of such studies. Moderately certain about the estimate of effect; further research may have an impact on our confidence in the estimate of effect and may change the estimate.

+++ – Marked effects.

Level of evidence – A: data derived from multiple randomised clinical trials or meta-analyses.

ACC, American College of Cardiology; AHA, American Heart Association; COR, Classification of recommendation; CVD, cardiovascular disease; DGAC, Dietary Guidelines Advisory Committee; EAS, European Atherosclerosis Society; ESC, European Society of Cardiology; LDL-C, low-density lipoprotein-cholesterol; MUFA, monounsaturated fatty acid; NHLBI, National Heart, Lung, and Blood Institute; PUFA, polyunsaturated fatty acid; RCT, randomised controlled trial; SFA, Saturated fatty acid.

events. In contrast, the recent AHA Presidential Advisory on dietary fats and CVD<sup>19</sup> concluded that ‘lowering intake of SFA and replacing it with unsaturated fats, especially PUFAs, will lower the incidence of CVD’. This conclusion was based on both epidemiological evidence as well as a meta-analysis of four well-controlled clinical trials, the latter of which demonstrated a 29% reduction in CHD events when SFA was replaced with PUFA. This disagreement reflects the studies included in the meta-analyses. Harcombe *et al*<sup>17</sup> and Hamley<sup>18</sup> included studies that were excluded by Sacks *et al*<sup>19</sup> due to methodological concerns (discussed in the AHA Presidential Advisory). Since this debate still continues, it is important to recognise the robust literature that demonstrates that lowering SFA

reduces LDL-cholesterol (table 2), an established causal factor in the development of atherosclerosis.<sup>20</sup>

### The need for rigorous dietary research: learning from the SFA debate

Dietary patterns are comprised of foods and nutrients, and this inter-relationship must be acknowledged in dietary research.<sup>21</sup> A reductionist approach has plagued SFA research and demonstrates the hazards of isolating a single nutrient and failing to account for the overall nutrient composition. This leads to confusion and unintended deleterious consequences (eg, a reduction in fat intake and an increase in refined carbohydrates) and impedes accurate translation of nutrition research.

**Table 2\***† Predicted effects of macronutrient replacement of dietary saturated fatty acids with PUFA, MUFA and carbohydrate on lipoprotein lipids based on results from controlled feeding trials\* Predicted effects\* on lipoprotein lipids of replacing 5% of energy from saturated fatty acids with the 5% of energy from the specified dietary component, mg/dL.

Dietary component	LDL-C	TG	HDL-C
PUFA	−9.0	−2.0	−1.0
MUFA	−6.5	+1.0	−6.0
CHO†	−6.0	+9.5	−2.0

Results are summarised from controlled feeding trials of subjects with average to mildly dyslipidaemic baseline levels of lipoprotein lipids. Effects may be more pronounced in those with higher baseline values. Taken from ref 4.

A meta-analysis of randomised controlled trials showed greater improvements in LDL-C with whole grains versus refined grains.<sup>23</sup> Compared with refined grains, whole grains reduced LDL-C by 3.48 mg/dL (95% CI 5.8 to 1.16;  $p < 0.01$ ). TG and HDL-C did not significantly improve in response to whole grains versus refined grains (TG mean difference −3.5 mg/dL, 95% CI −7.1 to 0.9,  $p = 0.10$ ; HDL-C mean difference −0.4, 95% CI −1.2 to 0.8,  $p = 0.59$ ).

CHO, Carbohydrate; HDL-C, high-density lipoprotein cholesterol; LDL-C, low-density lipoprotein-cholesterol; MUFA, monounsaturated fatty acid; PUFA, polyunsaturated fatty acid; TG, Triglycerides.

Future epidemiological research requires a more holistic approach to account for macronutrient substitutions. This translational approach favors support of healthy eating patterns and reduces unforeseen adverse effects more common when single nutrient recommendations are issued. Clinical research likewise benefits from dietary interventions that address the total dietary pattern including the influence of replacement foods and nutrients on the outcomes assessed.

### Summary and perspective

Although there is no biological requirement for SFA, a healthy diet will provide some SFA by virtue of meeting food-based dietary guidelines. A diet that limits SFA to <7% of calories accommodates a modest intake while achieving the recognised benefits of unsaturated fatty acids.

- Observational studies show cardiovascular benefits associated with lowering SFA and proportionally increasing PUFA, MUFA, whole grains and plant proteins.
- Epidemiological research that ignores nutrient differences in diets across the spectrum of SFA intake demonstrates no association of SFA with CVD risk. SFA intake is typically lowered and proportional increases in refined carbohydrates occur. Thus, the appropriate conclusion is that SFA and refined carbohydrates are equally deleterious on CVD risk. The controversy about the clinical trial evidence stems from studies with methodological problems and inclusion of these studies in meta-analyses.
- Recommendations to decrease SFA for CVD risk reduction are supported by robust evidence of a CVD benefit. The consensus of many scientists is to focus on a replacement message<sup>22 23</sup> that SFA be replaced with PUFA, MUFA, whole grains and plant proteins. This can be achieved with many healthy dietary patterns.
- Nutrition research must consider dietary patterns and the complex inter-relationship between dietary components to ensure translation into meaningful and health-promoting dietary recommendations.

**Contributors** All authors contributed to conceptualising and writing the manuscript.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient consent** Not required.

**Provenance and peer review** Commissioned; externally peer reviewed.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0>

### REFERENCES

- 1 Page IH, Stare FJ, Corcoran AC. Atherosclerosis and the fat content of the diet. *J Am Med Assoc* 1957;164:2048–51.
- 2 U.S. Department of Agriculture & U.S. Department of Health and Human Services. Dietary Guidelines for Americans. 1980. Available from: [https://www.cnpp.usda.gov/sites/default/files/dietary\\_guidelines\\_for\\_americans/1980thin.pdf](https://www.cnpp.usda.gov/sites/default/files/dietary_guidelines_for_americans/1980thin.pdf) [Accessed 12 Jul 2018].
- 3 Eckel RH, Jakicic JM, Ard JD, et al. American College of Cardiology/ American Heart Association Task Force on Practice Guidelines. 2013 AHA/ACC guideline on lifestyle management to reduce cardiovascular risk: a report of the American College of Cardiology/ American Heart Association Task Force on Practice Guidelines. *Circulation* 2014;129(25 Suppl 2):S76–99.
- 4 Jacobson TA, Maki KC, Orringer CE, et al. National lipid association recommendations for patient-centered management of dyslipidemia: part 2. *J Clin Lipidol* 2015;9(6 Suppl):S1–122.
- 5 U.S. Department of Health and Human Services, U.S. Department of Agriculture. Scientific report of the 2015 dietary guidelines advisory committee. Executive Summary. 2015. Available from: <https://health.gov/dietaryguidelines/2015-scientific-report/02-executive-summary.asp> [Accessed 12 Jul 2018].
- 6 Catapano AL, Graham I, De Backer G, et al. 2016 ESC/EAS Guidelines for the Management of Dyslipidaemias. *Eur Heart J* 2016;37:2999–3058.
- 7 Siri-Tarino PW, Sun Q, Hu FB, et al. Meta-analysis of prospective cohort studies evaluating the association of saturated fat with cardiovascular disease. *Am J Clin Nutr* 2010;91:535–46.
- 8 Stamler J. Diet-heart: a problematic revisit. *Am J Clin Nutr* 2010;91:497–9.
- 9 Katan MB, Brouwer IA, Clarke R, et al. Saturated fat and heart disease. *Am J Clin Nutr* 2010;92:459–60.
- 10 Pedersen JL, James PT, Brouwer IA, et al. The importance of reducing SFA to limit CHD. *Br J Nutr* 2011;106:961–3.
- 11 Chowdhury R, Warnakula S, Kunutsor S, et al. Association of dietary, circulating, and supplement fatty acids with coronary risk: a systematic review and meta-analysis. *Ann Intern Med* 2014;160:398–406.

- 12 de Souza RJ, Mente A, Maroleanu A, *et al.* Intake of saturated and trans unsaturated fatty acids and risk of all cause mortality, cardiovascular disease, and type 2 diabetes: systematic review and meta-analysis of observational studies. *BMJ* 2015;351:h3978.
- 13 Jakobsen MU, O'Reilly EJ, Heitmann BL, *et al.* Major types of dietary fat and risk of coronary heart disease: a pooled analysis of 11 cohort studies. *Am J Clin Nutr* 2009;89:1425–32.
- 14 Mozaffarian D, Micha R, Wallace S. Effects on coronary heart disease of increasing polyunsaturated fat in place of saturated fat: a systematic review and meta-analysis of randomized controlled trials. *PLoS Med* 2010;7:e1000252.
- 15 Li Y, Hruby A, Bernstein AM, *et al.* Saturated fats compared with unsaturated fats and sources of carbohydrates in relation to risk of coronary heart disease: a prospective cohort study. *J Am Coll Cardiol* 2015;66:1538–48.
- 16 Zong G, Li Y, Wanders AJ, *et al.* Intake of individual saturated fatty acids and risk of coronary heart disease in US men and women: two prospective longitudinal cohort studies. *BMJ* 2016;355:i5796.
- 17 Harcombe Z, Baker JS, DiNicolantonio JJ, *et al.* Evidence from randomised controlled trials does not support current dietary fat guidelines: a systematic review and meta-analysis. *Open Heart* 2016;3:e000409.
- 18 Hamley S. The effect of replacing saturated fat with mostly n-6 polyunsaturated fat on coronary heart disease: a meta-analysis of randomised controlled trials. *Nutr J* 2017;16:30.
- 19 Sacks FM, Lichtenstein AH, Wu JHY, *et al.* Dietary fats and cardiovascular disease: a presidential advisory From the American Heart Association. *Circulation* 2017;136:e1–e23.
- 20 Ference BA, Ginsberg HN, Graham I, *et al.* Low-density lipoproteins cause atherosclerotic cardiovascular disease. 1. Evidence from genetic, epidemiologic, and clinical studies. A consensus statement from the European Atherosclerosis Society Consensus Panel. *Eur Heart J* 2017;38:2459–72.
- 21 Tapsell LC, Neale EP, Satija A, *et al.* Foods, nutrients, and dietary patterns: interconnections and implications for dietary guidelines. *Adv Nutr* 2016;7:445–54.
- 22 Forouhi NG, Krauss RM, Taubes G, *et al.* Dietary fat and cardiometabolic health: evidence, controversies, and consensus for guidance. *BMJ* 2018;361:k2139.
- 23 Holl  nder PL, Ross AB, Kristensen M. Whole-grain and blood lipid changes in apparently healthy adults: a systematic review and meta-analysis of randomized controlled studies. *Am J Clin Nutr* 2015;102:556–72.