


Assessment of vitamin B₁₂ deficiency and B₁₂ screening trends for patients on metformin: a retrospective cohort case review

Darby Martin ¹, Jeet Thaker,¹ Maria Shreve,¹ Lois Lamerato,² Kartazyna Budzynska¹

To cite: Martin D, Thaker J, Shreve M, *et al.* Assessment of vitamin B₁₂ deficiency and B₁₂ screening trends for patients on metformin: a retrospective cohort case review. *BMJ Nutrition, Prevention & Health* 2021;4:e000193. doi:10.1136/bmjnp-2020-000193

¹Family Medicine, Henry Ford Hospital, Detroit, Michigan, USA
²Public Health Sciences, Henry Ford Health System, Detroit, Michigan, USA

Correspondence to

Dr Kartazyna Budzynska, Family Medicine, Henry Ford Hospital, Detroit, Michigan, USA; kbudzyn1@hfhs.org

Received 11 October 2020
Revised 12 December 2020
Accepted 17 December 2020
Published Online First
4 January 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

ABSTRACT

Objectives Our study investigated the use of vitamin B₁₂ testing in a large cohort of patients on metformin and assesses appropriateness and benefits of screening recommendations for vitamin B₁₂ deficiency.

Design This retrospective cohort study included insured adult patients who had more than 1 year of metformin use between 1 January 2010 and 1 October 2016 and who filled at least two consecutive prescriptions of metformin to establish compliance. The comparison group was not exposed to metformin. Primary outcome was incidence of B₁₂ deficiency diagnosed in patients on metformin. Secondary outcome was occurrence of B₁₂ testing in the patient population on metformin. Records dated through 31 December 2018 were analysed.

Setting Large hospital system consisting of inpatient and outpatient data base.

Participants A diverse, adult, insured population of patients who had more than 1 year of metformin use between 1 January 2010 and 1 October 2016 and who filled at least two consecutive prescriptions of metformin.

Results Of 13 489 patients on metformin, 6051 (44.9%) were tested for vitamin B₁₂ deficiency, of which 202 (3.3%) tested positive (vs 2.2% of comparisons). Average time to test was 990 days. Average time to test positive for deficiency was 1926 days. Factors associated with testing were linked to sex (female, 47.8%), older age (62.79% in patients over 80 years old), race (48.98% white) and causes of malabsorption (7.11%). Multivariable logistic regression showed older age as the only factor associated with vitamin B₁₂ deficiency, whereas African-American ethnicity approached significance as a protective factor.

Conclusions Based on our study's findings of vitamin B₁₂ deficiency in patients on metformin who are greater than 65 years old and have been using it for over 5 years, we recommend that physicians consider screening in these populations.

INTRODUCTION

Vitamin B₁₂ is a water-soluble vitamin. The human body extracts vitamin B₁₂ via gastric acid and intrinsic factor, which is subsequently absorbed in the terminal ileum.^{1 2} Patients who suffer from decreased ileal absorption are considered at increased risk of vitamin

B₁₂ deficiency.^{1 2} Other risk factors include decreased intrinsic factor production and decreased dietary intake.^{1 2} Medications including proton pump inhibitors, histamine H₂ blockers and metformin have also been shown to cause vitamin B₁₂ deficiency.^{1 2}

Although the exact mechanism is unknown, metformin's ability to cause vitamin B₁₂ deficiency is well established in the literature.³ The first reports describing this association were published in the late 1960s.⁴ A large meta-analysis of 19 intermediate and high-quality rated studies confirmed that patients on metformin will likely have lower vitamin B₁₂ levels.⁵ Another study showed that when the dose of metformin was increased by 100 mg, the odds for vitamin B₁₂ deficiency increased by 8%.⁶ Other studies reported similar findings.⁷ However, findings on the duration of metformin use and its effect on B₁₂ deficiency are inconsistent.⁶⁻⁹

The reported prevalence of vitamin B₁₂ deficiency from metformin ranges from 5.8% to 52%.^{5-7 10 11} Although no formal screening guidelines exist, the American Diabetes Association recommends that patients with long-term use of metformin, specifically those with anaemia or peripheral neuropathy, have vitamin B₁₂ levels measured periodically.¹² These recommendations are based on one of the largest and longest studies (over 2000 patients across 27 centres in the USA) reporting that patients on metformin, especially long-term use, are at higher risk for vitamin B₁₂ deficiency.¹³ The Agency for Healthcare Research and Quality recommends screening for vitamin B₁₂ deficiency in patients on long-term proton pump inhibitors or H₂ blockers (>12 months) and metformin (>4 months).¹⁴

In our study, we aimed to investigate use of vitamin B₁₂ testing in a large cohort of patients

on metformin and assess the benefit to formulating screening recommendations for vitamin B₁₂ deficiency.

METHODS

Study population

We received access to an administrative database of a non-profit health insurance plan offered by our hospital system to identify our system's insured adult patients (≥18 years) who had more than 1 year of metformin use between 1 January 2010 and 1 October 2016 and who filled at least two consecutive prescriptions of metformin. Using the criteria of filling at least two consecutive prescriptions of metformin and more than 1 year of metformin use, we assumed patient compliance to regularly taking the medication. Exclusion criteria included history of vitamin B₁₂ deficiency, a single prescription of metformin filled, and first metformin prescription filled within 30 days of insurance, as duration of metformin use could not be determined. Patients with only one prescription filled were deemed to be noncompliant with the medication and were excluded.

The comparison population included insured adult patients in the database within the same time frame who were not exposed to metformin.

Records dated through 31 December 2018 were analysed for vitamin B₁₂ testing and deficiency.

Measurements of metformin and outcome variables

Metformin use was determined from filled prescriptions by patients in the health plan database for the study period. We included all available doses, brands and combination pills of metformin. The main outcome variables were presence of testing for vitamin B₁₂ and vitamin B₁₂ deficiency defined as <180 pg/mL (determined by our institution laboratory department). Our institution uses Beckman Coulter DxI immunochemistry systems for vitamin B₁₂ testing.¹⁵

Data on additional covariates were collected from the health system's administrative databases, including demographic information (age, sex and race/ethnicity) and proton pump inhibitor use. Age was categorised into quintiles (18–39 years old, 40–49 years old, 50–64 years old, 65–79 years old and >80 years old based on population distribution). Confounder variables included malabsorption syndromes including small bowel disease, malnutrition, coeliac disease and previous bariatric surgery based on diagnosis and procedure codes in administrative data.

Screening guidelines

To be recommended for routine screening, a test such as vitamin B₁₂ must meet certain criteria based on Wilson and Jungner's principles of screening: (1) the disease is appropriate for screening (ie, the disease is serious, early treatment is beneficial and there is a high prevalence of preclinical disease); (2) screening is feasible and effective (eg, acceptable, cost-effective and detects many cases) and (3) the screening test is valid and widely available.^{16 17} We

kept these criteria in mind while performing our study and reviewed a random sample of charts to compare findings with these routine screening criteria.

Statistical analyses

Sample characteristics were described using mean and SD for continuous variables and frequencies (numbers and percentages) for categorical variables. Bivariate analysis was used to examine the effect of each of the covariates on incidence of testing and B₁₂ deficiency. Multivariable logistic regression models were used to examine the associations between metformin use and vitamin B₁₂ deficiency, adjusting for all other covariates. All analyses were performed using Epi Info V.7 (Centers for Disease Control and Prevention, Atlanta, Georgia).

RESULTS

Of 13 489 patients exposed to metformin, the majority were women (3461, 53.7%), aged 50–64 years (2316, 43.2%) to 65–79 years (2285, 31.1%) and African-American (2251, 40.1%) or white (2869, 43.4%) (table 1). Of the metformin exposed group, 6051 (44.9%) were tested for B₁₂ deficiency, of whom 202 (3.3%) were found to be B₁₂ deficient. The mean time between metformin initiation and B₁₂ testing was 972 days (the median time was 720 days). The mean time between metformin initiation and incidence of B₁₂ deficiency was 1926 days (5.3 years) (the median time was 2313 days).

Stratifying by demographics, men were tested for B₁₂ deficiency less often than women (41.5% vs 47.8%; OR 0.7568; 95% CI, 0.7047 to 0.8126; p<0.05) (table 2). Thus, men were tested approximately 25% less frequently compared with women. Men were also tested at a later date (mean=1037 days, median=785 days, positively skewed data set) compared with women (mean=922 days, median=675 days, positively skewed data set), which was also statistically significant (Kruskal-Wallis test, p<0.05). However, the vitamin B₁₂ deficiency rate in men (3.1%) compared with women (4.0%) was not statistically significant (table 3) (p=0.2031).

Analysis by age group showed that vitamin B₁₂ levels were evaluated at a much higher rate among the elderly (table 1). Patients aged 65–79 years and >80 years were tested for B₁₂ deficiency at a rate of 54.5% and 62.8%, respectively, higher than patients younger than 65 years old (<40% for ages 18–64 years). With each increase in age quintile, there was a 45% more likelihood of having B₁₂ levels evaluated (p<0.05). An older subgroup of patients (age >65 years) showed a B₁₂ deficiency rate of at least 4.2% compared with 2.5% in younger patients (p<0.05). With each increase in age quintile, there was approximately 34% more likelihood of incidence of B₁₂ deficiency (p<0.05). The elderly (age >80 years) were tested much sooner after starting metformin than the study population (mean of 785 days vs 970 days) and were found to have B₁₂ deficiency much sooner than the study population (mean of 1491 days vs 1926 days).

Table 1 Characteristics of the population and associated testing

| | Total no | Ever tested | Mean days to test | Median days to test | Deficiency | Mean days to B ₁₂ deficiency | Median days to B ₁₂ deficiency |
|---|--------------|--------------|-------------------|---------------------|------------|---|---|
| Gender | | | | | | | |
| Male | 6247 (46.3%) | 2590 (41.5%) | 1037 | 785 | 81 (3.1%) | 1883 | 1921 |
| Female | 7242 (53.7%) | 3461 (47.8%) | 922 | 675 | 121 (3.5%) | 1955 | 2407 |
| Total | 13489 (100%) | 6051 (44.9%) | 972 | 720 | 202 (3.3%) | 1926 | 2313 |
| Age | | | | | | | |
| 18–39 | 830 (6.2%) | 292 (35.2%) | 929 | 643 | 3 (1.0%) | 2686 | 2780 |
| 40–49 | 1716 (12.7%) | 581 (33.9%) | 942 | 695 | 12 (2.1%) | 2367 | 2690 |
| 50–64 | 5828 (43.2%) | 2316 (39.7%) | 1050 | 780 | 64 (2.8%) | 2019 | 2397 |
| 65–79 | 4196 (31.1%) | 2285 (54.5%) | 952 | 723 | 99 (4.3%) | 1894 | 2076 |
| >80 | 919 (6.8%) | 577 (62.8%) | 785 | 553 | 24 (4.2%) | 1492 | 1133 |
| Total | 13489 (100%) | 6051 (44.9%) | 972 | 720 | 202 (3.3%) | 1926 | 2313 |
| Race | | | | | | | |
| African-American | 5409 (40.1%) | 2251 (41.6%) | 966 | 724 | 59 (2.6%) | 2132 | 2407 |
| Asian | 499 (3.7%) | 209 (41.9%) | 905 | 684 | 6 (2.9%) | 2312 | 2680 |
| Hispanic | 338 (2.5%) | 149 (44.1%) | 1184 | 791 | 6 (4.0%) | 2230 | 2500 |
| Other/unknown | 1385 (10.3%) | 573 (41.4%) | 969 | 692 | 22 (3.8%) | 1756 | 2165 |
| Caucasian | 5858 (43.4%) | 2869 (49.0%) | 970 | 723 | 109 (3.8%) | 1811 | 2040 |
| Total | 13489 (11%) | 6051 (44.9%) | 972 | 720 | 202 (3.3%) | 1926 | 2313 |
| Confounders | | | | | | | |
| Malabsorption disorders | 694 (5.1%) | 498 (71.8%) | 860 | 572 | 11 (2.2%) | 1864 | 2046 |
| Proton pump inhibitor use (prescriptions) | 3957 (29.3%) | 2173 (54.9%) | 947 | 703 | 70 (3.2%) | 2058 | 2452 |
| Total | 4651 (100%) | 2671 (57.4%) | 972 | 720 | 81 (3.0) | 1926 | 2313 |

Analysis by race showed no significant difference in rates of testing, rates of B₁₂ deficiency, days between metformin initiation and B₁₂ testing or days between metformin initiation and detection of B₁₂ deficiency. However, the African-American population was tested at a lower rate (41.62%) compared with the non-African-American population (44.9%), showing a 15% less likelihood of being tested compared with the other ethnic groups ($p<0.05$). The incidence of B₁₂ deficiency in the African-American group was 27% lower than other races, which approached statistical significance ($p=0.051$) (table 3).

Evaluation of patients with malabsorption disorders who were started on metformin showed a much higher B₁₂ testing rate than the entire study population (70% vs 44.86%). Patients with malabsorption disorders were 3.64 times more likely to be tested than patients without malabsorption disorders ($p<0.01$). The incidence of vitamin B₁₂ deficiency in this group, however, was not significantly lower than the total study population (2.2% vs 3.34%, $p=0.38$). Patients using proton pump inhibitors while on metformin were 64% more likely to be tested for vitamin B₁₂ deficiency ($p<0.05$); however, the B₁₂ deficiency rate was not statistically significant.

Table 2 Logistic regression analysis: ever tested for B₁₂ deficiency

| | OR | 95% CI | P value |
|---------------------------|------|----------------|------------------|
| Age | 1.46 | 1.402 to 1.514 | <0.001 |
| AfricanAmerican | 0.85 | 0.793 to 0.918 | <0.001 |
| Malabsorption syndromes | 3.64 | 3.064 to 4.334 | <0.001 |
| Male | 0.76 | 0.705 to 0.813 | <0.001 |
| Proton pump inhibitor use | 1.64 | 1.519 to 1.771 | <0.001 |

Bold values are statistically significant.

Table 3 Logistic regression analysis: B₁₂ deficiency

| | OR | 95% CI | P value |
|---------------------------|------|----------------|------------------|
| African-American | 0.73 | 0.534 to 1.001 | 0.05 |
| Age | 1.34 | 1.141 to 1.573 | <0.001 |
| Male | 0.83 | 0.619 to 1.107 | NS |
| Malabsorption syndromes | 0.76 | 0.407 to 1.409 | NS |
| Proton pump inhibitor use | 0.92 | 0.681 to 1.233 | NS |

Bold values are statistically significant.
NS, not statistically significant.

Of the 202 charts indicating B₁₂ deficiency, we reviewed a random sample of 105 charts to identify the most common reasons for vitamin B₁₂ testing in our study population. Anaemia (26%), cognitive decline (17%) and neuropathy (16%) were the most commonly reported reasons, whereas only 3.8% stated that the B₁₂ level was ordered for concern of deficiency caused by metformin.

DISCUSSION

In our study population, individuals on metformin were tested for vitamin B₁₂ deficiency more frequently if they were elderly, had a malabsorption disorder, or were taking proton pump inhibitors. African-American patients were tested less frequently compared with other races, and men were tested less frequently compared with women. The only factor associated with vitamin B₁₂ deficiency, however, was older age. African-American ethnicity approached significance as a protective factor for B₁₂ deficiency.

While we found testing to be suboptimal, the incidence of vitamin B₁₂ deficiency for patients on metformin was lower than we had expected (3.3% vs 2.2% comparison). Our findings are concordant with the Diabetes Prevention Programme Outcomes Study (DPPOS) (B₁₂ deficiency rate of 4.3% vs placebo deficiency rate of 2.4%).^{13 18} The DPPOS study population consisted of 68% women, 45% from ethnic and racial minority groups, and 20% aged 60 years or older.^{13 18} Our study consisted of 53% women and 56.6% minorities. The DPPOS did not distinguish between the different minority races in their report.^{13 18} The DPPOS study defined B₁₂ deficiency as <203 pg/mL, a higher value compared with our definition.^{13 18} Another large study that used data from the National Health and Nutrition Examination Survey (NHANES) also showed similar B₁₂ deficiency results.¹⁹ Their study had 49.4% women, but few minorities (14.6% African-American and 10.9% Hispanic).¹⁹ Their definition of B₁₂ deficiency was <148 pmol/L, a lower value compared with our definition.¹⁹ The rate of B₁₂ deficiency of the NHANES study population without diabetes was similar to ours.¹⁹

Our study's racial composition was different from other large studies. With our population including 40% African-Americans, our results showed that this ethnicity was less likely to be deficient in B₁₂. The US population includes 13.4% African-Americans.²⁰ The NHANES and DPPOS studies did not have large African-American populations but found similar results regarding their lower B₁₂ deficiency rates. This finding suggests a possibility that the African American race may be protective from vitamin B₁₂ deficiency, but more research is needed to explore this hypothesis, as the African-American group was tested at lower rates. This could be a factor that influenced the diagnosis of B₁₂ deficiency, as, historically, African-Americans have remained undiagnosed in other cases.

Our study showed that patients using metformin for more than 5 years were at greater risk for vitamin B₁₂ deficiency. The study using NHANES data found that

the likelihood of vitamin B₁₂ deficiency was greater the longer patients took metformin (B₁₂ deficiency rate of 4.1% when on metformin for 3–10 years vs 8.1% for >10 years).¹⁹ The DPPOS study found that patients on metformin had a similar B₁₂ deficiency prevalence at 5-year follow-up compared with patients in the NHANES study.¹³ Given that our results are similar to both studies, we agree that longer use of metformin may increase the likelihood of B₁₂ deficiency.

We also found increased age to be correlated with vitamin B₁₂ deficiency in our study. The NHANES study found the rate of B₁₂ deficiency to increase by a factor of 1 with each year of age¹⁹ whereas the DPPOS study found no relation between age and B₁₂ deficiency.¹³ This is a particularly important point as studies have shown that metformin can be useful in the treatment of cancers, obesity, liver diseases, renal diseases and cardiovascular diseases which tend to plague older populations.²¹

We used results from our study to analyse whether the 1.2% difference in vitamin B₁₂ deficiency rates between our metformin and comparison populations was significant enough to indicate the need for screening for B₁₂ deficiency. Using criteria for creating a screening guideline, not all can be met. Vitamin B₁₂ deficiency can cause many symptoms that can be reversed with vitamin B₁₂ supplementation. Supplementation is not expensive; an injection at our health system is approximately US\$10 and is usually covered by insurance. The test for B₁₂ deficiency, however, is not affordable at this time. Patients at our institution are charged US\$80 per B₁₂ test, which would only be covered if associated with a symptom of deficiency as predetermined by the patient's insurance. As indicated by a random sample chart review, 59% of our patients had symptoms of B₁₂ deficiency (eg, anaemia, neuropathy or cognitive decline) that led to testing vs true screening. This correlates with the current recommendations.

Our study is unique in that we were able to distinguish the rates at which testing occurred and the rates at which different populations were tested. Neither the DPPOS nor NHANES study reported these rates. We found that women were tested more than men, older patients were tested more frequently compared with their younger counterparts, and African-Americans were less likely to be tested. Many factors account for the testing habits of physicians based on these populations' characteristics. More research regarding this topic is necessary.

Our study has several limitations. Due to the retrospective design, we were not able to completely address all potential confounders such as dietary habits, nutritional status, metformin use and vitamin B₁₂ supplementation. We used data that were coded in our electronic medical system. Our study did not stratify based on metformin dosing or duration of metformin prescription (30 days vs 90 days). Our patients were insured under a premium insurance plan that includes few Medicare patients and no Medicaid patients. This limits the generalisability of the study and could potentially lead to selection bias, but helped us confirm that the patient filled some scripts of

metformin. Metformin compliance could not be verified, although we excluded patients with less than two filled metformin prescriptions and those who filled a metformin prescription within 30 days of starting their insurance. Nutritional status was not accounted for, including over-the-counter supplements and vegan and vegetarian diets. We defined vitamin B₁₂ deficiency as <180 pg/mL whereas other studies used <203 pg/mL, which makes study comparisons more difficult.

Despite its limitations, our study accounted for confounders and excluded preexisting vitamin B₁₂ deficiency to assess incidence and not prevalence. We used a single insurance database to confirm that metformin prescriptions were filled by our patients and to estimate duration of metformin use via documentation of the first and subsequent filled prescriptions. Our study population was diverse and overall included more minorities than white patients.

CONCLUSION

Our study findings suggest that physicians should be cognizant of the increased incidence of vitamin B₁₂ deficiency in select populations. These populations include patients with greater than 5 years of metformin use and age greater than 65 years old. The patient's race should also be considered.

What this paper adds

- ▶ Question: Should we screen for vitamin B₁₂ deficiency in patients taking metformin?
- ▶ Findings: Patients who are greater than 65 years old and patients who have been taking metformin for more than 5 years are at increased risk. The black population is also likely at increased risk.
- ▶ Meaning: Consider screening for B₁₂ deficiency in these populations even if they are asymptomatic for the deficiency.

Acknowledgements We thank the residents Pooja Kulkarni, MD, Islam Zin El Din, MD, Rehab Bakir, MD, Diahann Marshall, MD, Berta Rezik, MD and Andrea Smith, DO, for their assistance with the study. We thank Roger Tuttleman for their help with the data analysis.

Contributors DM, JT, LL and KB set up the data protocol, performed chart review, analysed and interpreted the data and took the lead in the writing process. MS made substantial contributions to the conception and design, and actively participated in writing manuscript and interpreting data. DM is the guarantor of this work and, as such, had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. All authors read and approved the final manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval This retrospective cohort study was approved by our health system's institutional review board with waiver of signed consent based on use of existing administrative data.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data may be obtained from a third party and are not publicly available. Deidentified participant data requests will be reviewed on

request by The Henry Ford Health System Regulatory Counsel. Please contact the corresponding author for more information.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID ID

Darby Martin <http://orcid.org/0000-0001-6025-4182>

REFERENCES

- 1 Langan RC, Goodbred AJ. Vitamin B12 deficiency: recognition and management. *Am Fam Physician* 2017;96:384–9.
- 2 Hunt A, Harrington D, Robinson S. Vitamin B12 deficiency. *BMJ* 2014;349:g5226.
- 3 de Jager J, Kooy A, Leher P, et al. Long term treatment with metformin in patients with type 2 diabetes and risk of vitamin B-12 deficiency: randomised placebo controlled trial. *BMJ* 2010;340:c2181.
- 4 Tomkin GH, Hadden DR, Weaver JA, et al. Vitamin-B12 status of patients on long-term metformin therapy. *Br Med J* 1971;2:685–7.
- 5 Chapman LE, Darling AL, Brown JE. Association between metformin and vitamin B₁₂ deficiency in patients with type 2 diabetes: A systematic review and meta-analysis. *Diabetes Metab* 2016;42:316–27.
- 6 de Groot-Kamphuis DM, van Dijk PR, Groenier KH, et al. Vitamin B12 deficiency and the lack of its consequences in type 2 diabetes patients using metformin. *Neth J Med* 2013;71:386–90.
- 7 Beulens JWJ, Hart HE, Kuijs R, et al. Influence of duration and dose of metformin on cobalamin deficiency in type 2 diabetes patients using metformin. *Acta Diabetol* 2015;52:47–53.
- 8 Ko S-H, Ko S-H, Ahn Y-B, et al. Association of vitamin B12 deficiency and metformin use in patients with type 2 diabetes. *J Korean Med Sci* 2014;29:965–72.
- 9 Haeusler S, Parry-Strong A, Krebs JD. The prevalence of low vitamin B12 status in people with type 2 diabetes receiving metformin therapy in New Zealand—a clinical audit. *N Z Med J* 2014;127:8–16.
- 10 Rodríguez-Gutiérrez R, Montes-Villarreal J, Rodríguez-Velver KV, et al. Metformin use and vitamin B12 deficiency: Untangling the association. *Am J Med Sci* 2017;354:165–71.
- 11 Pflipsen MC, Oh RC, Sagui A, et al. The prevalence of vitamin B(12) deficiency in patients with type 2 diabetes: a cross-sectional study. *J Am Board Fam Med* 2009;22:528–34.
- 12 American Diabetes Association. Standards of medical care in diabetes-2018. *Diabetes Care* 2018;41:S1–159.
- 13 Aroda VR, Edelstein SL, Goldberg RB, et al. Long-Term metformin use and vitamin B12 deficiency in the diabetes prevention program outcomes study. *J Clin Endocrinol Metab* 2016;101:1754–61.
- 14 Agency for Healthcare Research and Quality. Guideline summary: Cobalamin (vitamin B12) deficiency - investigation & management. Available: <https://www2.gov.bc.ca/assets/gov/health/practitioner-pro/bc-guidelines/cobalamin.pdf> [Accessed 18 Apr 2019].
- 15 Beckman Coulter, Inc. Vitamin B12 assay procedure. Available: <https://www.beckmancoulter.com/products/immunoassay/access-active-b12#/documents> [Accessed 19 Nov 2020].
- 16 Dobrow MJ, Hagens V, Chafe R, et al. Consolidated principles for screening based on a systematic review and consensus process. *CMAJ* 2018;190:E422–9.
- 17 Wilson JMG, Jungner G. *Principles and practice of screening for disease*. Geneva: World Health Organization, 1968.
- 18 Diabetes Prevention Program Research Group, Knowler WC, Fowler SE, et al. 10-Year follow-up of diabetes incidence and weight

- loss in the diabetes prevention program outcomes study. *Lancet* 2009;374:1677–86.
- 19 Reinstatler L, Qi YP, Williamson RS, *et al.* Association of biochemical B₁₂ deficiency with metformin therapy and vitamin B₁₂ supplements: the National health and nutrition examination survey, 1999–2006. *Diabetes Care* 2012;35:327–33.
- 20 United States Census Bureau. Population estimates- race and Hispanic origin. Available: www.census.gov/quickfacts/fact/table/US/PST045218#PST045218 [Accessed 18 Apr 2019].
- 21 Lv Z, Guo Y. Metformin and its benefits for various diseases. *Front Endocrinol* 2020;11:191. <https://doi.org/10.3389/fendo.2020.00191>.