Vitamin C and alcohol: a call to action

Daniel James Lim,¹ Yogesh Sharma,² Campbell Henry Thompson³¹Adelaide Medical School,
University of Adelaide, Adelaide,
South Australia, Australia²Department of General
Medicine, Flinders Medical
Centre, Bedford Park, South
Australia, Australia³Discipline of Medicine,
University of Adelaide, Adelaide,
South Australia, Australia

Correspondence to

Daniel James Lim, Adelaide
Medical School, University of
Adelaide, Adelaide, SA 5005,
Australia; danlimjre@gmail.com

Received 25 September 2018

Revised 24 October 2018

Accepted 1 November 2018

Published Online First

5 December 2018

INTRODUCTION

Vitamin C, an essential water-soluble nutrient that cannot be synthesised by humans, has a role in numerous biological reactions.¹ It is necessary for enzymatic reactions and biosynthesis of hormones,² and protects biomolecules through its antioxidant properties.¹ Critical in the biosynthesis of collagen, it is crucial in preserving essential tissue structure and function, without which capillary fragility develops.^{1,2}

While its biological function has been well proven, its therapeutic utility has been well contested over the years. The initial hope of a positive association between vitamin C status and health from several large cohort trials^{3–5} was quashed by results from large randomised

controlled trials (RCTs) that did not find significant positive effects of vitamin C supplementation, when mortality or morbidity were set as endpoints.^{6–8} A review by Lykkesfeldt and Poulsen⁹ remarked this to be a result of a non-specific approach to examining the isolated effect of vitamin C, and lamented a lack of subpopulation definition in the trials. However, the subsequent shift in the public's and health authorities' interests has left a gaping hole in evidence regarding vitamin C replacement.

Chronic alcoholism, on the other hand, remains a significant and widespread public health issue, whereby a recent global study found alcohol to be the seventh leading risk factor for death and disability-adjusted life-years.¹⁰

Table 1 Case reports on vitamin C deficiency and chronic alcohol intake

Reference	Country	Sex	Age	Clinical signs	Alcohol intake (drinks/day)	Plasma vitamin C (µmol/L) (RR 28–70)	Other deficiencies	Replacement regimen		
								Vitamin C/day (mg)	Route	Period
Chaudhry <i>et al</i> ¹¹	England	Male	44	Lethargy, FH, PH, G	Chronic	<3	NS	800	Oral	2 weeks
Wang and Still ¹²	USA	Male	65	E, PH, G, CH	10–15	6.82	NS	1000	NS	5 days
								500	NS	After 5 days
Léger ¹³	Canada	Male	47	Lethargy, E, PH	8–10	<10	NS	NS	Oral	NS
Takeshima <i>et al</i> ¹⁴	Japan	Male	45	P, G, Oe	13	1.13	Vit B ₉ : 1.3 ng/mL (RR 2.4–12.0) Hb: 58 g/L (>135)	1200	Oral	3 weeks
Gião Antunes <i>et al</i> ¹⁵	Portugal	Male	40	GH, FH, PO	12	7.95	Vit B ₉ : 2.1 ng/mL (RR 2.4–12.0) Hb: 71 g/L (>135)	300	Oral	2 weeks
								100	Oral	1 month after 2 weeks
Lux-Battistelli and Battistelli ¹⁶	USA	Male	54	Lethargy, weak, P, GH	Dependent	3.1	NS	1000	Oral	8–10 weeks
		Male	24	Fatigue, weak, dyspnoea, Oe	14–28	<3	NS	1000	Oral	12 weeks
		Male	48	FH, unsteady	34	<24	NS	500	Oral	12 weeks
Bennett <i>et al</i> ¹⁷	USA	Male	64	E, GH, unsteady weight loss	8–10	5.68	Vit B ₉ : 2 ng/mL (RR >4.7) Vit B ₆ : <2 µg/L (RR 5–50)	NS	Oral	NS

CH, corkscrew hairs; E, ecchymoses; FH, follicular hyperkeratosis; G, gingivitis; GH, gingival hypertrophy; Hb, haemoglobin; NS, not specified; Oe, oedema; P, petechiae or purpura; PH, perifollicular haemorrhages; PO, periodontitis; RR, reference range; Vit, vitamin.

© Author(s) (or their employer(s)) 2018. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Lim DJ, Sharma Y, Thompson CH. *bmjnph* 2018;1:17–22.

There has been no lack of case reports documenting florid scurvy in chronic alcoholics,^{11–17} yet the mechanistic links between hypovitaminosis C and chronic alcoholism remain postulations based on research from the early 1980s. Furthermore, replacement regimens and routes remain far from standardised, as evidenced by the varied treatment approaches in these reports (table 1).

As such, the present review examines the current and past knowledge on vitamin C deficiency and its role in chronic alcoholics, evaluates the current guidelines on alcohol withdrawal, and provides suggestions and exhortation for future trials. PubMed, MEDLINE and Current Contents (1960–2018) were searched using medical subject headings and free text with a combination of the following terms: vitamin C, vitamin C deficiency, ascorbic acid, scurvy, alcohol, chronic alcoholic and alcohol withdrawal guidelines. Original articles, contemporary reviews, case reports and international guidelines were used.

Clinical significance of vitamin C deficiency

The prevalence of hypovitaminosis varies across the world; however, it has been observed to be higher in hospitalised patients.¹⁸ The current definition of optimal vitamin C status, while controversial, has been based on a dietary intake that results in a plasma concentration of approximately 70 µmol/L in an otherwise healthy person.¹⁹ The recommended daily allowance for vitamin C has been stated at 75 mg/day for women and 90 mg/day for men in the USA¹⁹; however, its uptake is highly dose-dependent and varies depending on saturation from daily diet.

Furthermore, the extensive variability of plasma vitamin C levels to clinical signs has prompted the definition of hypovitaminosis C with arbitrary cut-offs. Severe vitamin C deficiency has been proposed to be at a serum level of <11 µmol/L,²⁰ while marginal vitamin C deficiency is suggested as serum levels of >11 µmol/L to 28 µmol/L.²⁰

Scurvy, the clinical hallmark of severe vitamin C deficiency, is often the red flag that alerts the clinician to measure plasma vitamin C levels. It manifests primarily as capillary haemorrhages, gingivitis, lassitude and muscle weakness.^{11–17} Apart from scurvy, severe vitamin C deficiency has been shown to increase the relative risk of myocardial infarction,⁵ while marginal vitamin C deficiency has also been shown to be significantly associated with all-cause mortality, with weak associations with death from cancer.⁴

Despite the significant progress being made in the clinical significance of hypovitaminosis C, little is known about its role or replacement in chronic alcoholics.

The case for vitamin C deficiency and alcohol

Excessive alcohol consumption (>80 g/day) has been identified as one of the many risk factors for scurvy.¹⁸ This has traditionally been thought to be due to malnutrition with poor living conditions, leading to multiple nutritional deficiencies, especially vitamin B₁ (thiamine) deficiency, associated with Wernicke-Korsakoff syndrome,

Figure 1 Postulated mechanisms of chronic alcohol intake and vitamin C replacement. ¹Majumdar *et al.*²² — — —, Prevents; ETOH, ethanol/alcohol.

and vitamin B₃ (niacin) deficiency, associated with pellagra.^{11–17} However, as early as 1978, in a cohort of 35 patients with alcohol-related illness, Baines²¹ reported the incidence of vitamin C deficiency to be 91%, compared with that of vitamin B₁ deficiency reported at 31%. Since then, studies from the 1980s have shown that hypovitaminosis C alone is likely to have a significant effect on chronic alcohol use and vice versa. However, there has since been a lack of follow-up on these key findings that may play a significant role in the detoxification or withdrawal regimen of the chronic alcoholic.

In 1981, Majumdar *et al.*²² found that short-term intravenous vitamin C supplementation (500 mg mane for 5 days) in chronic alcoholics with hypovitaminosis C improved serum levels of vitamin C significantly, but in 16 of 25 patients they did not return to normal ranges. This was consistent with previous studies, where it took up to 3 months of oral vitamin C supplementation for blood levels to return to normal in chronic alcoholics.²³

Thus, Majumdar and colleagues²² postulated that ethanol-induced enterocyte toxicity induces intestinal malabsorption of vitamins, while ethanol-induced hepatotoxicity results in insufficient hepatic transformation of many vitamins to their active metabolites²² (figure 1). They also suggested that vitamin C may be protective against toxic effects of acetaldehyde, as proven in animal studies.²⁴ This may reduce hepatotoxicity and possibly the biochemical basis of addiction, given acetaldehyde's action in dopaminergic stimulation of opiate receptors in the brain.²² However, given the low number of patients (n=25) in their study, the likelihood of insufficient power,

Figure 2 Study findings of acute alcohol intake and vitamin C. ¹Faizallah *et al.*²⁶ ²Susick and Zannoni.²⁷ ETOH, ethanol/ alcohol.

and therefore a false-negative finding, was high, requiring further large trials to confirm such results.

Since then, research looking at isolated vitamin C replacement in chronic alcoholism has been sporadic, barring a recent breakthrough showing decreased vitamin C uptake in pancreatic acinar cells of chronic alcoholics.²⁵

The relationship between acute alcohol consumption and vitamin C deficiency has received more interest during the same time period (figure 2). Alcohol consumption of up to 0.58 g ethanol/kg of body weight as whisky/lager was shown, in 1986, to produce a 47% increase in urinary ascorbic acid excretion, compared with controls who drank nothing.²⁶ Thereafter, a robust study by Susick and Zannoni²⁷ in 1987 showed that vitamin C pretreatment (5 g daily for 2 weeks) significantly enhanced blood ethanol clearance, suggesting a contributory role of vitamin C to ethanol oxidation. These findings were bolstered by Chen *et al.*²⁸ in 1990 showing significantly increased plasma alcohol clearance following both short-term and long-term vitamin C pretreatment.

Susick and Zannoni's²⁷ study also looked at behavioural measures, finding that pretreatment resulted in significantly less ethanol-induced impairment of motor coordination and colour discrimination by >20%, as compared with placebo.²⁷ These findings support a neurochemical role for the vitamin, as previously suggested by Majumdar *et al.*²²

While these studies were important, the populations looked at were healthy, unlike those of chronic alcohol use.

Present-day guidelines and attitudes

Given the strong bidirectional, and possibly therapeutic, link between vitamin C treatment and alcohol

consumption, one would have expected a push for trials looking at vitamin C supplementation in chronic alcoholics. However, to date, vitamin C supplementation has only been proven to ameliorate ethanol-induced hepatotoxicity in mice,²⁹ not humans, while RCTs, as stated above,^{6–8} have not looked into the at-risk subpopulation of chronic alcoholic use. This is manifested in the present-day guidelines on alcohol withdrawal which provide scant instruction on vitamin C supplementation regimen or route (table 2).

The WHO guidelines for withdrawal state in a single line 'Multivitamin supplements and particularly vitamin B1 (thiamine) supplements (at least 100 mg daily during withdrawal) should also be provided',³⁰ while yet another international guideline, published in 2017 by the World Federation of Societies of Biological Psychiatry, makes mention of the state of hypovitaminosis in the alcoholic, but none to vitamin C or its replacement.³¹

Guidelines from multiple task forces across the United States of America (USA) make no mention at all of vitamin C replacement despite a known 'state of hypovitaminosis' in alcoholics,^{32–33} while Australian guidelines provide brief instruction, such as 'oral multivitamin preparation can be given to nutritionally depleted patients for several days'; however, there is no graded recommendation nor evaluation of the evidence.^{34–35}

While in the United Kingdom (UK), although national guidelines do not address hypovitaminosis C,³⁶ local guidelines of various National Health Service Foundation Trusts^{37–39} both identify hypovitaminosis and specifically mention intravenous replacement of vitamin B and C acutely, although in the setting of Wernicke's encephalopathy. Encouragingly, they mention parenteral vitamin supplementation acutely, followed by oral supplementation. The lack of specificity, however, given that vitamin C is delivered adjunctively in an ampoule combined with vitamin B, makes it difficult to investigate the isolated therapeutic value of vitamin C replacement.

Given the lack of evidence, promising RCT findings or even trials regarding vitamin C supplementation in chronic alcoholics, it comes as no surprise that major guidelines to date have not looked beyond the specifics of vitamin B₁ and B₁₂ replacement regimens. It seems the once-gilded promise of vitamin C's antioxidant value has lost its shine.

Yet it may be judicious to heed recommendations of past studies. Majumdar *et al.*²² on the basis of their findings, advised nearly 40 years ago to include prolonged replenishment of vitamin C in conventional detoxification therapy for ethanol withdrawal syndrome, preferably by the intravenous route, postulating that it is needed for the utilisation of other vitamins and nutrients as well.²² Today, we know this to be true, given that vitamin C, with its reducing and chelating properties, is known to be the most efficient enhancer of non-haem iron absorption.⁴⁰

Given the immense therapeutic potential of vitamin C replacement and robust evidence from past studies, it seems that the current practice gap or guideline omission

Table 2 International guidelines on alcohol withdrawal and vitamin C replacement

Country	Organisation	Recommendation				
		Alerts to hypovitaminosis	Vitamin C replacement	Route	Dose	Time period
International	WHO ³⁰	+	–	–	–	–
	World Federation of Societies of Biological Psychiatry ³¹	+	–	–	–	–
USA	American Psychiatric Association ³²	–	–	–	–	–
	Substance Abuse and Mental Health Services ³³	+	Multivitamin	Intravenous	NS	NS
Australia	Australian Government, Department of Health and Ageing ³⁴	+	Multivitamin	Oral	Daily	NS
	Government of South Australia, SA Health ³⁵	+	Multivitamin	NS	NS	NS
UK	National Institute for Health and Care Excellence ³⁶	–	–	–	–	–
	Southern Health, NHS Foundation Trust ³⁷	+	Vitamins B and C*	Intramuscular	1 Amp daily	5 days
			Multivitamin	Oral	1 Tab daily	6 weeks
	Guy's and St Thomas', NHS Foundation Trust ³⁸	+	Vitamins B and C*	Intravenous	1–2 Amp three times a day	7 days
	NHS Grampian ³⁹	+	Vitamins B and C*	Intravenous	1–2 Amp three times a day	5 days
				Oral	500 mg	NS

*Given as Pabrinex injection.

(+), present; (–), absent; Amp, ampoule (5 mL each); NS, not specified; Tab, tablet.

may stem from a non-targeted approach to the subpopulation of chronic alcoholics.

Difficulties faced in populations with chronic alcohol use and recommendations for future studies

There are multiple complexities faced when designing trials and studies for chronic alcoholics. Malnutrition, and its subsequent polyvitamin deficiency, is rampant in populations of chronic alcohol use,²² making it challenging to confer directionality to mechanistic hypotheses or isolate the effect of an intervention due to a single substance.

Given the association of inflammation and vitamin C deficiency, due to an increased tissue expenditure of nutrients,¹⁸ the systemic inflammatory response seen in chronic alcoholics with liver cirrhosis only adds another layer of complexity to any analysis. The subsequent loss of liver synthetic function also confers difficulty in interpreting whether clinical signs, such as widespread ecchymoses, are a result of a lack of vitamin C or coagulation factors, or both.^{11–17} In addition, progressive muscle weakness and gait unsteadiness in scurvy can mimic ethanol-induced cerebellar toxicity.

There also remain conflicting views concerning the necessity of vitamin C supplementation. A recent review by Flannery *et al*,⁴¹ on the management of alcohol-associated vitamin and electrolyte deficiencies, suggested providing routine supplementation of high-dose thiamine, magnesium sulfate and folic acid. They did not, however, recommend routine supplementation with vitamin C, citing a

lack of published data investigating the efficacy or safety of the acute use of multivitamin injection in patients with possible alcohol withdrawal. Despite this, they did remark that ‘this lack of data does not necessarily negate the potential benefits of intravenous multivitamin in the acute setting’.⁴¹

After reviewing 35 RCTs with vitamin C as an intervention, Lykkesfeldt and Poulsen⁹ made the following suggestions for study parameters: hypovitaminosis C as an entry-level inclusion, verification of vitamin C levels at entry and during the study by a validated method, adjustment for confounders, a valid hypothesis or molecular mechanism involving vitamin C or mechanism-related hard clinical endpoint as a primary outcome, and explicit exclusion criteria.⁹

Targeting a subpopulation of chronic alcoholics, we suggest stratification of the quantity of alcohol intake and classification of severity of alcoholic liver disease, and adjusting for smoking status, socioeconomic status, dietary intake, quality of life, cognitive assessment, depression scale and acute illness, as vitamin C deficiency has been associated with these parameters.^{11–17}

We suggest instituting vitamin C replacement intravenously with a specified regimen. This may be difficult to justify given the current lack of evidence for intravenous replacement, while juggling known risks, such as cannula site infection and hyperoxaluria, one of the side effects of high-dose vitamin C replacement.⁴²

We therefore hope this review will serve as a call for more research, in particular for future RCTs to be designed with an inclusion criterion of hypovitaminosis C in a population affected by chronic alcohol use. The endpoints of such trials should include haematological, dermatological and hepatological clinical and laboratory measures (table 1) as surrogate markers for primary or secondary outcomes. This would provide the necessary evidence to further explore the optimal route and regimen of vitamin C replacement. Additionally, more observational studies are needed to evaluate the risk factors for hypovitaminosis C and its likelihood in patients of chronic alcohol exposure.

CONCLUSION

In conclusion, we have found a significant absence in the literature and published guidelines concerning vitamin C deficiency and chronic alcoholics. Given the substantial public health burden of chronic alcoholism, and the relatively inexpensive means of vitamin C replacement, we would like to call for an urgent response to the 'dire need to examine the effect of vitamin C as a single supplement in populations which have been carefully defined', as Lykkesfeldt and Poulsen⁹ so aptly adjudged.⁹ There is a need to build on, and to not forget, the insightful and robust evidence from the 1980s–1990s when developing trials and studies exploring the replacement of vitamin C deficiency in the chronic alcoholic, in order to review and further bolster current guidelines.

Contributors DJL engaged in the literature search, data collection for figures and tables, data analysis and interpretation, and supervised manuscript writing and revisions. YS analysed and interpreted the data and provided suggestions in drafting the manuscript. CHT engaged in the analysis and interpretation of data, data collection for figures and tables, and manuscript writing and revision.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0>

REFERENCES

- Englund S, Seifter S. The biochemical functions of ascorbic acid. *Annu Rev Nutr* 1986;6:365–406.
- May JM, Qu ZC, Meredith ME. Mechanisms of ascorbic acid stimulation of norepinephrine synthesis in neuronal cells. *Biochem Biophys Res Commun* 2012;426:148–52.
- Khaw KT, Bingham S, Welch A, et al. Relation between plasma ascorbic acid and mortality in men and women in EPIC-Norfolk prospective study: a prospective population study. European Prospective Investigation into Cancer and Nutrition. *Lancet* 2001;357:657–63.
- Loria CM, Klag MJ, Caulfield LE, et al. Vitamin C status and mortality in US adults. *Am J Clin Nutr* 2000;72:139–45.
- Nyyssönen K, Parviainen MT, Salonen R, et al. Vitamin C deficiency and risk of myocardial infarction: prospective population study of men from eastern Finland. *BMJ* 1997;314:634–8.
- Cook NR, Albert CM, Gaziano JM, et al. A randomized factorial trial of vitamins C and E and beta carotene in the secondary prevention of cardiovascular events in women: results from the Women's Antioxidant Cardiovascular Study. *Arch Intern Med* 2007;167:1610–8.
- Heart Protection Study Collaborative Group. MRC/BHF Heart Protection Study of antioxidant vitamin supplementation in 20,536 high-risk individuals: a randomised placebo-controlled trial. *Lancet* 2002;360:23–33.
- Hercberg S, Galan P, Preziosi P, et al. The SU.VI.MAX Study: a randomized, placebo-controlled trial of the health effects of antioxidant vitamins and minerals. *Arch Intern Med* 2004;164:2335–42.
- Lykkesfeldt J, Poulsen HE. Is vitamin C supplementation beneficial? Lessons learned from randomised controlled trials. *Br J Nutr* 2010;103:1251–9.
- Griswold MG, Fullman N, Hawley C, et al. Alcohol use and burden for 195 countries and territories, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *The Lancet* 2018;392:1015–35.
- Chaudhry SI, Newell EL, Lewis RR, et al. Scurvy: a forgotten disease. *Clin Exp Dermatol* 2005;30:735–6.
- Wang AH, Still C. Old world meets modern: a case report of scurvy. *Nutr Clin Pract* 2007;22:445–8.
- Léger D. Scurvy: reemergence of nutritional deficiencies. *Can Fam Physician* 2008;54:1403–6.
- Takeshima M, Echizenya M, Inomata Y, et al. Scurvy in an alcohol-dependent patient with a severely unbalanced diet. *Psychiatry Clin Neurosci* 2014;68:242–3.
- Gião Antunes AS, Peixe B, Guerreiro H. Gastrointestinal bleeding secondary to scurvy in an alcoholic malnourished cirrhotic patient. *ACG Case Rep J* 2017;4:e29.
- Lux-Battistelli C, Battistelli D. Latent scurvy with tiredness and leg pain in alcoholics: an underestimated disease three case reports. *Medicine* 2017;96:e8861.
- Bennett SE, Schmitt WP, Stanford FC, et al. Case 22-2018: A 64-year-old man with progressive leg weakness, recurrent falls, and anemia. *N Engl J Med* 2018;379:282–9.
- Fain O, Pariés J, Jacquart B, et al. Hypovitaminosis C in hospitalized patients. *Eur J Intern Med* 2003;14:419–25.
- Institute of Medicine & National Academy of Sciences. *Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium and Carotenoids. A Report of the Panel on Dietary Antioxidants and Related Compounds, Subcommittees on Upper Reference Levels of Nutrients and of the Interpretation and Use of Dietary Reference Intakes, and the Standing Committee on the Scientific Evaluation of Dietary Reference Intakes, Food and Nutrition Board*. Washington, DC: National Academy Press, 2000.
- Johnston CS, Corte C. People with marginal vitamin C status are at high risk of developing vitamin C deficiency. *J Am Diet Assoc* 1999;99:854–6.
- Baines M. Detection and incidence of B and C vitamin deficiency in alcohol-related illness. *Ann Clin Biochem* 1978;15:307–12.
- Majumdar SK, Patel S, Shaw GK, et al. Vitamin C utilization status in chronic alcoholic patients after short-term intravenous therapy. *Int J Vitam Nutr Res* 1981;51:274–8.
- Sviripa EV. [Vitamin C metabolism in alcoholism and alcoholic psychoses]. *Zh Nevropatol Psikhiatr Im S S Korsakova* 1971;71:422–5.
- Sprince H, Parker CM, Smith GG. Article on Ascorbic Acid. *Int J Vitam Nutr Res* 1977;(Suppl 16):185.
- Subramanian VS, Srinivasan P, Said HM. Uptake of ascorbic acid by pancreatic acinar cells is negatively impacted by chronic alcohol exposure. *Am J Physiol Cell Physiol* 2016;311:C129–35.
- Faizallah R, Morris AI, Krasner N, et al. Alcohol enhances vitamin C excretion in the urine. *Alcohol Alcohol* 1986;21:81–4.
- Susick RL, Zannoni VG. Effect of ascorbic acid on the consequences of acute alcohol consumption in humans. *Clin Pharmacol Ther* 1987;41:502–9.
- Chen MF, Boyce HW, Hsu JM. Effect of ascorbic acid on plasma alcohol clearance. *J Am Coll Nutr* 1990;9:185–9.
- Li Wenjie, Xin Qiliang, Xin Qiliang, et al. Vitamin C protective role for alcoholic liver disease in mice through regulating iron metabolism. *Toxicol Ind Health* 2011;27:341–8.
- World Health Organization. *Clinical Guidelines for Withdrawal Management and Treatment of Drug Dependence in Closed Settings. WHO Guidelines Approved by the Guidelines Review Committee*. Geneva: WHO Press, 2009.

31. Soyka M, Kranzler HR, Hesselbrock V, et al. Guidelines for biological treatment of substance use and related disorders, part 1: Alcoholism, first revision. *World J Biol Psychiatry* 2017;18:86–119.
32. Herbert K, Roger W, Raymond A. *Practice Guideline For The Treatment of Patients With Substance Use Disorders*. 2nd edn. American Psychiatric Association [Internet], 2006.
33. Center for Substance Abuse Treatment. *Detoxification and Substance Abuse Treatment. Treatment Improvement Protocol (TIP) Series 45. DHHS Publication No. (SMA) 06-4131*. Rockville, MD: Substance Abuse and Mental Health Services Administration, 2006.
34. Australian Government Department of Health and Ageing. *Guidelines for the Treatment of Alcohol Problems*. Sydney: Commonwealth of Australia, 2009: 1–232.
35. Drug and Alcohol Services South Australia. *Management of patients at risk of alcohol withdrawal in acute hospitals*. Adelaide: Government of South Australia Department for Health and Ageing, 2016: 8.
36. National Institute for Health and Clinical Excellence, 2018. Alcohol-use disorders: diagnosis and clinical management of alcohol-related physical complications (CG100). Available from: <https://www.nice.org.uk/guidance/CG100> [Accessed 24 Aug 2018].
37. National Health Service. *Alcohol Withdrawal Guidelines*. Southampton: Southern Health NHS Foundation Trust, 2018: 7–10.
38. National Health Service. *Clinical Guideline: Detection of alcohol misusers, management of alcohol withdrawal syndrome and Wernicke's Encephalopathy*. 2. London: Guy's and St Thomas' NHS Foundation Trust, 2012.
39. National Health Service. *NHS grampian (acute sector) guidance for the prescribing of vitamin and mineral supplements for hospitalized patients with a history of excess alcohol consumption*. Aberdeen: NHS Grampian, 2013: 3–6.
40. Teucher B, Olivares M, Cori H. Enhancers of iron absorption: ascorbic acid and other organic acids. *Int J Vitam Nutr Res* 2004;74:403–19.
41. Flannery AH, Adkins DA, Cook AM. Unpeeling the evidence for the Banana Bag: Evidence-based recommendations for the management of alcohol-associated vitamin and electrolyte deficiencies in the ICU. *Crit Care Med* 2016;44:1545–52.
42. Baxmann AC, De O G Mendonça C, Heilberg IP. Effect of vitamin C supplements on urinary oxalate and pH in calcium stone-forming patients. *Kidney Int* 2003;63:1066–71.