Vitamin D and SARS-CoV-2 virus/ COVID-19 disease

Susan A Lanham-New,¹ Ann R Webb,² Kevin D Cashman,³ Judy L Buttriss,⁴ Joanne L Fallowfield,⁵ Tash Masud,⁶ Martin Hewison,⁷ John C Mathers,⁸ Mairead Kiely,³ Ailsa A Welch,⁹ Kate A Ward,¹⁰ Pamela Magee,¹¹ Andrea L Darling,¹ Tom R Hill,⁸ Carolyn Greig,¹² Colin P Smith,¹³ Richard Murphy,¹⁴ Sarah Leyland,¹⁵ Roger Bouillon,¹⁶ Sumantra Ray,^{11,17,18} Martin Kohlmeier ^{18,19}

To cite: Lanham-New SA, Webb AR, Cashman KD, *et al.* Vitamin D and SARS-CoV-2 virus/COVID-19 disease. *BMJ Nutrition, Prevention & Health* 2020;3:e000089. doi:10.1136/bmjnp-2020-000089

For numbered affiliations see end of article.

Correspondence to

Professor Susan A Lanham-New, Nutritional Sciences, University of Surrey Faculty of Health and Medical Sciences, Guildford GU2 7XH, UK; s.lanham-new@surrey.ac.uk

Received 16 April 2020
Revised 29 April 2020
Accepted 30 April 2020
Published Online First
13 May 2020

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

BACKGROUND AND AIM

The spread of novel SARS-CoV-2 virus, and the disease COVID-19 that is caused by SARS-CoV-2, continues apace. Saving lives and slowing the worldwide pandemic remain of utmost importance to everyone: the public, healthcare professionals, scientists, industry and governments.

It is absolutely essential that advice given to the public is evidence-based, accurate and timely; anything less would mislead and has the potential to cause harm. Popular information channels, such as social media platforms, have been rife with misinformation that has been perpetuated by fear and uncertainty. This has been the case particularly for diet and lifestyle advice. There are recommendations for the prevention of the spread of COVID-19 from the WHO,¹ the UK,² Irish³ and USA⁴ governments and the European Commission,⁵ as well as public health and healthcare agencies, including key direction on self-isolation.⁶

This short original report aims to provide a balanced scientific view on vitamin D and SARS-CoV-2 virus/COVID-19 disease. It provides a succinct summary of the current scientific evidence of associations between vitamin D, influenza, upper respiratory tract infections (URTIs) and immune health. Importantly, the paper concludes with lifestyle strategies for avoiding vitamin D deficiency and ensuring a healthy balanced diet at any time, including during the current pandemic. The overarching messages are as follows: (1) Vitamin D is essential for good health. (2) Many people, particularly those living in northern latitudes (such as the UK, Ireland, Northern Europe, Canada and the northern parts of the USA, northern India and China), have poor vitamin D status, especially in winter or if confined indoors. (3) Low

vitamin D status may be exacerbated during this COVID-19 crisis (eg, due to indoor living and hence reduced sun exposure), and anyone who is self-isolating with limited access to sunlight is advised to take a vitamin D supplement according to their government's recommendations for the general population (ie, 400 IU/day for the UK⁷ and 600 IU/day for the USA (800 IU for >70 years))⁸ and the European Union (EU).⁹ (4) There is no strong scientific evidence to show that very high intakes (ie, mega supplements) of vitamin D will be beneficial in preventing or treating COVID-19. (5) There are evidenced health risks with excessive vitamin D intakes especially for those with other health issues such as a reduced kidney function.

NUTRITION, VITAMIN D AND IMMUNITY

Good nutrition and lifestyle factors (such as physical activity) have a positive impact on immune function, promoting biological and physiological systems and processes that enable humans to resist infection. In light of the current COVID-19 pandemic, and given the importance of diet to overall health and well-being, nutrients (macro and micro) deserve special attention.¹⁰ As a key micronutrient, vitamin D should be given particular focus—not as a ‘magic bullet’ to beat COVID-19, as the scientific evidence base is severely lacking at this time—but rather as part of a healthy lifestyle strategy to ensure that populations are nutritionally in the best possible place.¹¹

Vitamin D is unique: it is a prohormone which is produced in the skin during exposure to sunlight (UVB radiation at 290–315 nm) with, usually, smaller amounts obtained from food. During the winter months in areas of middle-high latitude, the solar elevation

remains low throughout the short daylight period, and there is insufficient solar UVB to support appreciable vitamin D synthesis.¹² For most people, dietary intake does not fully supply the body's vitamin D needs and so vitamin D status declines during the winter. For example, in Manchester, UK (53.5N) the nadir of seasonal vitamin D status occurs in February, with sunlight exposure once again becoming effective for vitamin D synthesis in the skin only from March onwards.¹³ Relatively high prevalence of low vitamin D status globally has been reported over recent decades in a wide range of population groups,¹⁴ including those in low latitude areas (despite the abundance of sunlight) and not necessarily confined to winter.¹⁵ This may be due to environmental factors, such as air pollution, as well as cultural factors that lead to skin being covered and not subject to sunlight exposure.¹⁶ Older, housebound individuals are at particularly high risk of vitamin D deficiency.¹⁷

Vitamin D status is reflected by the level of the circulating metabolite 25-hydroxyvitamin D (25OHD), which is produced by hepatic hydroxylation of vitamin D coming from either skin or the gut from oral intake.¹⁸ If the 25OHD concentration is low (as defined in the UK by a 25OHD concentration of <25 nmol/L⁷ and in the USA and some other countries by a 25OHD concentration of <30 nmol/L),^{8,9} such as observed commonly during and towards the end of the winter, this indicates that stores are depleted and vitamin D-requiring functions may be impaired. The association between low vitamin D status and increased risk of rickets in children and poor musculoskeletal health in adults is well documented. In addition, vitamin D, via its active metabolites, regulates more than 200 genes including those genes that are responsible for cellular proliferation, differentiation and apoptosis.¹⁹ The discovery of the expression of nuclear vitamin D receptors and vitamin D metabolic enzymes in immune cells provides a scientific rationale for the potential role of vitamin D in maintaining immune homeostasis and in preventing the development of autoimmune processes.²⁰ The field of vitamin D research has grown exponentially in recent years with a much improved understanding of its biological importance. Recent meta-analyses of randomised controlled trials (RCTs) concluded that the use of vitamin D supplements was associated with lower total mortality in elderly, mostly vitamin D-deficient participants.^{21,22} However, the most recent Vitamin D Assessment Study and the Vitamin D and Omega-3 Fatty Acid Study did not show a mortality effect in vitamin D-replete adults.^{23,24}

VITAMIN D AND RESPIRATORY HEALTH

It has been hypothesised that there is an association between seasonal URTIs and low vitamin D status because both occur in the winter months. However, controversy remains as to whether there is a direct link between the seasonality of influenza and vitamin D deficiency. Higher influenza incidence in winter may be due to behavioural

reasons including the greater time spent indoors, which increases individuals' proximity and hence likely interpersonal transmission. Nonetheless, vitamin D appears to inhibit pulmonary inflammatory responses²⁵ while enhancing innate defence mechanisms against respiratory pathogens.²⁶ Moreover, population-based studies show positive associations between circulating 25OHD concentration and lung function.²⁷ Nevertheless, formal systematic reviews/meta-analyses of these associations are urgently required.

URTIs, or 'common colds', are the most widespread of infectious diseases, with more than 200 viruses contributing to the clinical symptoms. Epidemiological studies in children have found a strong association between URTI and rickets.²⁸ A large cross-sectional study of the US population reported that URTI infections were higher in those with lower vitamin D status, with the association being stronger in those with respiratory diseases such as asthma and chronic obstructive pulmonary disease.²⁹ There is evidence that lower vitamin D status is associated with acute respiratory tract infections (ARTIs).³⁰ In a recent systematic review and meta-analysis of individual participant data from vitamin D supplementation RCTs, vitamin D supplementation reduced the risk of ARTI, with the greatest benefit in those with vitamin D deficiency at baseline.³¹ However, it is important to note the limitations to this systematic review/meta-analysis;^{32,33} there was a high level of heterogeneity in the findings and concomitantly, the overall significant results in the meta-analysis of the 24 included trials was dependent on the inclusion of the two studies undertaken in developing countries: Mongolia and Afghanistan. These two trials had specific participants and the findings should not be extrapolated to populations from more developed countries. This is particularly important as Panagiotou *et al* (2013) demonstrated that the efficacy of many treatments was substantially greater in less developed compared with more developed countries.³⁴ Furthermore, the specific clinical definitions of ARTI were varied across included studies, with many research participants with ARTI being self-diagnosed.

VITAMIN D, SARS-COV-2 VIRUS AND COVID-19 DISEASE

The continued spread of the novel SARS-CoV-2 virus, and the disease COVID-19 that is caused by SARS-CoV-2, has led to calls for widespread high-dose vitamin D supplementation.³⁵⁻³⁹ These calls are without support from pertinent studies in humans at this time, but rather based on speculations about *presumed* mechanisms. There have been two key studies published to support this presumption: (1) An unbiased screen of repurposed drugs for treatment of avian influenza A H5N1 virus using appropriate cell lines and mice, which highlighted calcitriol (the active hormone of vitamin D) as a potential therapy.⁴⁰ (2) A recent analysis of vitamin D and viral infections.⁴¹ However, whether these mechanisms apply with SARS-CoV-2 is not known. Studies investigating

vitamin D and COVID-19 are currently underway,^{42 43} and more are likely to follow. Given that ethnic minorities are disproportionately affected with Covid-19—and this appears to be the case principally in the UK, the USA and other European countries—further research is justified, especially given that there is clear evidence that vitamin D deficiency is particularly common in these ethnic groups.^{44 45}

However, we strongly caution against doses higher than the upper limit (4000 IU/day; 100 µg/day),^{7 8} and certainly of very high doses of vitamin D (in some reports, 10 000 IU/day (250 µg/day) of vitamin D are being promoted) unless under personal medical advice/clinical advice by a qualified health professional. Instead, we advocate the following lifestyle strategies for avoiding vitamin D deficiency and ensuring a healthy, balanced diet.

VITAMIN D ADVICE

Current evidence-based advice for the prevention of vitamin D deficiency includes:

1. *Supplementation with vitamin D according to Government guidelines* (eg, 400 IU/day (10 µg/day) for the UK;⁷ 600 IU/day (15 µg/day) for the USA⁸ (800 IU/day (20 µg/day) for >70 years) and Europe.⁹ These recommendations were established to ensure that 25OHD concentrations in the majority of the population are above 25 nmol/L (UK) in order to protect musculoskeletal health or above 30 nmol/L (USA) to minimise the risk of vitamin D deficiency (the USA recommendation was also established to optimise musculoskeletal health in the population using a 25OHD concentration of 50 nmol/L). Supplementation with vitamin D is particularly important during times of self-isolation associated with limited sunlight exposure. This is in line with the UK Scientific Advisory Committee on Nutrition (SACN)⁷ recommendations for vitamin D, and the US Institute of Medicine (IOM) recommendations for vitamin D,⁸ both of which were established under the assumption of minimal exposure to sunlight. Thus, re-emphasis of advice on safe sun exposure (below) and reinforcing government advice on supplements especially when sunlight exposure is low would further boost vitamin D status. The UK SACN, US IOM and EU European Food Safety Agency recommend that vitamin D intake (total from both foods and dietary supplements) should be limited to 4000 IU/day (100 µg/day) for adults, and there is broad international consensus that the general public should avoid higher dose supplements that risk total intake from all sources exceeding this level.
2. *Consumption of a nutritionally balanced diet*, for example, according to the UK Eatwell Guide⁴⁶ and US Food Pyramid⁴⁷ including vitamin D rich foods, that is, oily fish, red meat, egg yolk and fortified foods,⁴⁸ such as breakfast cereals in the UK, as well as fortified milk in the USA and Canada.^{49 50}

3. *Safe sunlight exposure to boost vitamin D status*. Safe sunlight exposure will enable vitamin D production in skin from March through September in the UK, and at most northern latitudes. Dermal synthesis of vitamin D is most efficient with short, regular (daily) exposures when the sun is at its strongest (in the middle of the day).⁵¹ The efficiency of vitamin D synthesis declines well before the threshold for sunburn is reached¹² but the desirable dose is skin-type dependent and so exposure times required differ for different skin types.⁵² For the UK about 10 min of exposure at around lunch-time, in-season appropriate clothing, can meet vitamin D needs for white-skinned people; this increases to about 25 min for those of skin type V (ie, South Asian, brown skin tones).^{53 54} What is key is to try to achieve the sunlight exposure without leaving home (eg, in the garden/balcony); and if that is not possible ensure that social distancing is maintained at all times. Increasing the unprotected skin area (skin not protected by clothing or sunscreen) will increase the vitamin D supply from skin while keeping exposure times short and suberythematous. Exposing as much skin as temperature and social comfort allow will maximise vitamin D supply through this route. For those of skin type V and VI (brown or black skin) the exposure requirements in UK sunlight are more challenging to achieve than for white-skinned people and oral vitamin D intake is especially important.
4. *Appropriate diet and lifestyle measures*, as emphasised by the WHO at this time, including adequate nutrition to protect the immune system.⁵⁵
5. *Targeted nutritional advice*, for example, for UK Military personnel as advised by the Defence Nutrition Advisory Service, with specific reference to COVID-19.⁵⁶
6. *Vitamin D—advice for bone health*. The Royal Osteoporosis Society provides specific guidelines on the management of vitamin D deficiency in adults with, or at risk of developing, bone disease.⁵⁷

In conclusion, we recommend appropriate vitamin D RCTs to evaluate the effects of vitamin D supplementation on COVID-19 infections. Until there is more robust scientific evidence for vitamin D, we strongly caution against the use of high vitamin D supplementation (greater than the upper limit of 4000 IU/day (100 µg/day)). Rather, we strongly endorse avoidance of vitamin D deficiency⁵⁸ in the population (as per the six points above) and complete adherence to government's advice worldwide on the *prevention of the spread of COVID-19*.

SUMMARY FOR SOCIAL MEDIA

Vitamin D is essential for good health, especially bone and muscle health. Many people have low blood levels of vitamin D, especially in winter or if confined indoors, because summer sunshine is the main source of vitamin D for most people. Government vitamin D intake recommendations for the general population are 400 IU (10 µg) per day for the UK⁷ and 600 IU (15 µg) per day for the

USA⁸ (800 IU (20 µg) per day for >70 years) and the EU.⁹ Taking a daily supplement (400 IU /day (10 µg/day) in the UK) and eating foods that provide vitamin D is particularly important for those self-isolating⁵⁹ with limited exposure to sunlight. Vitamin D intakes greater than the upper limit of 4000 IU (100 µg) per day may be harmful and should be avoided unless under personal medical/clinical advice by a qualified health professional.

Author affiliations

¹Department of Nutritional Sciences, School of Biosciences and Medicine, Faculty of Health & Medical Sciences, University of Surrey, Guildford, UK

²Earth and Environmental Sciences, University of Manchester, Manchester, UK

³School of Food and Nutritional Sciences, University College Cork, Cork, Ireland

⁴British Nutrition Foundation, London, UK

⁵Applied Physiology, Institute of Naval Medicine, Gosport, UK

⁶Department of Medicine, Nottingham University Hospitals NHS Trust, Nottingham, UK

⁷Institute of Metabolism and Systems Research, University of Birmingham, Birmingham, UK

⁸Human Nutrition Research Centre, Population Health Sciences Institute, Newcastle University, Newcastle, UK

⁹Norwich Medical School, University of East Anglia, Norwich, UK

¹⁰MRC Lifecourse Epidemiology, NIHR Biomedical Research Centre, University of Southampton and University Hospitals Southampton NHS Foundation Trust, Southampton, UK

¹¹School of Biomedical Sciences, Ulster University at Coleraine, Coleraine, UK

¹²School of Sport, Exercise and Rehabilitation Sciences and MRC-Versus Arthritis Centre for Musculoskeletal Ageing Research, University of Birmingham, Birmingham, UK

¹³School of Pharmacy and Biomolecular Sciences, University of Brighton, Brighton, UK

¹⁴Centre for Environment & Sustainability, Faculty of Engineering & Physical Sciences, University of Surrey, Guildford, UK

¹⁵Royal Osteoporosis Society, Camerton, Bath, UK

¹⁶Laboratory of Clinical and Experimental Endocrinology, Department of Chronic Diseases, Metabolism and Ageing, KU Leuven, 3000 Leuven, Belgium

¹⁷School of Humanities and Social Sciences, University of Cambridge, Cambridge, UK

¹⁸NNEEdPro Global Centre for Nutrition and Health, St John's Innovation Centre, Cambridge, UK

¹⁹School of Medicine & Gillings School of Global Public Health, University of North Carolina at Chapel Hill & UNC Nutrition Research Institute, Chapel Hill, North Carolina, USA

Twitter Kate A Ward @kateaward17

Contributors All authors drafted and amended the vitamin D and SARS-CoV-2 virus/COVID-19 disease report and all were in agreement with the wording. SAL-N, ARW and KDC are collective lead authors.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None

Patient consent for publication Not required.

Ethics approval Not required

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Martin Kohlmeier <http://orcid.org/0000-0003-1478-1744>

REFERENCES

- World Health Organisation. Coronavirus disease (COVID-19) advice for the public, 2020. Available: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/advice-for-public>
- UK Government. Coronavirus (COVID-19): guidance, 2020. Available: <https://www.gov.uk/government/collections/coronavirus-covid-19-list-of-guidance>
- Irish Government. COVID-19 (Coronavirus), 2020. Available: <https://www.gov.ie/en/campaigns/c36c85-covid-19-coronavirus/>
- US Government. Government response to coronavirus, COVID-19, 2020. Available: <https://www.usa.gov/coronavirus>
- EU Commission. Overview of the Commission's response, 2020. Available: https://ec.europa.eu/info/live-work-travel-eu/health/coronavirus-response/overview-commissions-response_en
- NHS. What to do if you or someone you live with has symptoms of coronavirus, 2020. Available: <https://www.nhs.uk/conditions/coronavirus-covid-19/self-isolation-advice/>
- Scientific Advisory Committee on Nutrition. Vitamin D and health, 2016. Available: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/537616/SACN_Vitamin_D_and_Health_report.pdf
- Institute of Medicine. *Dietary reference intakes for calcium and vitamin D*. Washington, DC, USA, 2011.
- European Food Safety Agency. Dietary reference values for vitamin D, 2016. Available: <http://www.efsa.europa.eu/en/efsajournal/pub/4547>
- Calder PC, Carr AC, Gombart AF, et al. Optimal nutritional status for a well-functioning immune is an important factor to protect against viral infections, 2020. Available: <https://www.preprints.org/manuscript/202003.0199/v1>
- Hewison M. Vitamin D and immune function: an overview. *Proc Nutr Soc* 2012;71:50–61.
- Webb AR, Kline L, Holick MF. Influence of season and latitude on the cutaneous synthesis of vitamin D3: exposure to winter sunlight in Boston and Edmonton will not promote vitamin D3 synthesis in human skin. *J Clin Endocrinol Metab* 1988;67:373–8.
- Rhodes LE, Webb AR, Fraser HI, et al. Recommended summer sunlight exposure levels can produce sufficient (> or =20 ng ml⁻¹) but not the proposed optimal (> or =32 ng ml⁻¹) 25(OH)D levels at UK latitudes. *J Invest Dermatol* 2010;130:1411–8.
- Bouillon R. Vitamin D status is worse in Africa than in other continents. *Lancet* 2019;8:e20–1.
- Mendes MM, Charlton K, Thakur S, et al. Future perspectives in addressing the global issue of vitamin D deficiency. *Proc Nutr Soc* 2020;79:246–51.
- Al-Yatama FI, AlOtaibi F, Al-Bader MD, et al. The effect of clothing on vitamin D status, bone turnover markers, and bone mineral density in young Kuwaiti females. *Int J Endocrinol* 2019;2019:6794837.
- Griffin TP, Wall D, Blake L, et al. Vitamin D status of adults in the community, in outpatient clinics, in hospital and in nursing homes in the West of Ireland. *J Gerontol A Biol Sci Med Sci* 2020. doi:10.1093/gerona/glaa010. [Epub ahead of print: 14 Jan 2020].
- Bikle DD. Vitamin D metabolism, mechanism of action, and clinical applications. *Chem Biol* 2014;21:319–29.
- Umar M, Sastry KS, Chouchane AI. Role of vitamin D beyond the skeletal function: a review of the molecular and clinical studies. *Int J Mol Sci* 2018;19. doi:10.3390/ijms19061618. [Epub ahead of print: 30 May 2018].
- Vanherwegen A-S, Gysemans C, Mathieu C. Vitamin D endocrinology on the cross-road between immunity and metabolism. *Mol Cell Endocrinol* 2017;453:52–67.
- Bjelakovic G, Gluud LL, Nikolova D, et al. Vitamin D supplementation for prevention of mortality in adults. *Cochrane Database Syst Rev* 2014;1:CD007470.
- Keum N, Lee DH, Greenwood DC, et al. Vitamin D supplementation and total cancer incidence and mortality: a meta-analysis of randomized controlled trials. *Ann Oncol* 2019;30:733–43.
- Scragg R. The Vitamin D Assessment (ViDA) study - Design and main findings. *J Steroid Biochem Mol Biol* 2020;198:105562.
- Manson JE, Cook NR, Lee I-M, et al. Vitamin D supplements and prevention of cancer and cardiovascular disease. *N Engl J Med* 2019;380:33–44.
- Hughes DA, Norton R. Vitamin D and respiratory health. *Clin Exp Immunol* 2009;158:20–5.
- Zdrenghea MT, Makrinioti H, Bagacean C, et al. Vitamin D modulation of innate immune responses to respiratory viral infections. *Rev Med Virol* 2017;27:1–3.
- Craveiro V, Cabral M, Araújo J, et al. Association of serum 25-hydroxyvitamin D concentration with pulmonary function in young adults. *Nutrients* 2018;10. doi:10.3390/nu10111728. [Epub ahead of print: 11 Nov 2018].

- 28 Patwari A, Nabi G, Nadroo AM, *et al.* Pulmonary changes in rickets in children. *Indian Pediatr* 1979;16:413–5.
- 29 Ginde AA, Mansbach JM, Camargo CA. Association between serum 25-hydroxyvitamin D level and upper respiratory tract infection in the third National health and nutrition examination survey. *Arch Intern Med* 2009;169:384–90.
- 30 Laaksi I, Ruohola J-P, Tuohimaa P, *et al.* An association of serum vitamin D concentrations < 40 nmol/L with acute respiratory tract infection in young Finnish men. *Am J Clin Nutr* 2007;86:714–7.
- 31 Martineau AR, Jolliffe DA, Hooper RL, *et al.* Vitamin D supplementation to prevent acute respiratory tract infections: systematic review and meta-analysis of individual participant data. *BMJ* 2017;356:i6583.
- 32 Hemilä H. Statistical problems in the vitamin D and respiratory infection meta-analysis. *BMJ* 2018;356:i658.
- 33 McIndoe KS. Vitamin D supplementation to prevent acute respiratory tract infections. *BMJ* 2017;356.
- 34 Panagiotou OA, Contopoulos-Ioannidis DG, Ioannidis JPA. Comparative effect sizes in randomised trials from less developed and more developed countries: meta-epidemiological assessment. *BMJ* 2013;346:f707.
- 35 Grant WB, Lahore H, McDonnell SL, *et al.* Evidence that vitamin D supplementation could reduce risk of influenza and COVID-19 infections and deaths. *Nutrients* 2020;12:E988.
- 36 McCartney DM, Byrne DG. Optimisation of vitamin D status for enhanced Immuno-protection against Covid-19. *Ir Med J* 2020;113:58.
- 37 Garami AR. Preventing a covid-19 pandemic - is there a magic bullet to save COVID-19 patients? We can give it a try! *BMJ Comments*. *BMJ* 2020;368.
- 38 Schwalfenberg GK. Preventing a COVID-19 pandemic. rapid response. *BMJ* 2020;368.
- 39 Vitamin D Wiki. Coronavirus can likely be fought by vitamin D., 2020. Available: <https://vitamindwiki.com/COVID19+Coronavirus+can+most+likely+be+fought+by+Vitamin+D>
- 40 Huang F, Zhang C, Liu Q, *et al.* Identification of amitriptyline HCl, flavin adenine dinucleotide, azacitidine and calcitriol as repurposing drugs for influenza A H5N1 virus-induced lung injury. *PLoS Pathog* 2020;16:e1008341.
- 41 Lee C. Controversial Effects of vitamin D and related genes on viral infections, pathogenesis, and treatment outcomes. *Nutrients* 2020;12:962.
- 42 Chinese Clinical Trials Registry. The relationship between vitamin D and novel coronavirus pneumonia (COVID-19), 2020. Available: <http://www.chictr.org.cn/showprojen.aspx?proj=51390>
- 43 Chinese Clinical Trials Registry. Impact of vitamin D deficiency on prognosis of patients with novel coronavirus pneumonia (COVID-19), 2020. Available: <http://www.chictr.org.cn/showprojen.aspx?proj=49302>
- 44 Darling AL, Hart KH, Macdonald HM, *et al.* Vitamin D deficiency in UK South Asian women of childbearing age: a comparative longitudinal investigation with UK Caucasian women. *Osteoporos Int* 2013;24:477–88.
- 45 Kramer H, Camacho P, Aloia J, *et al.* Association between 25-hydroxyvitamin D and intact parathyroid hormone levels across latitude among adults with African ancestry. *Endocr Pract* 2016;22:911–9.
- 46 Public Health England. The Eatwell guide, 2016. Available: <https://www.gov.uk/government/publications/the-eatwell-guide>
- 47 US Department of Health and Human Services. USA food pyramid, 2020. Available: <https://health.gov/our-work/food-nutrition>
- 48 British Nutrition Foundation. 8 tips for eating well, 2018. Available: <https://www.nutrition.org.uk/healthyliving/healthydiet/8tips.html>
- 49 Buttriss JA, Lanham-New SA. Is a vitamin D fortification strategy needed? in press. *Nutrition Bulletin* 2020.
- 50 National Institute of Health. Vitamin D, 2020. Available: <https://ods.od.nih.gov/factsheets/VitaminD-health%20Professional/>
- 51 Webb AR, Engelsen O. Calculated ultraviolet exposure levels for a healthy vitamin D status. *Photochem Photobiol* 2006;82:1697–703.
- 52 Webb AR. Who, what, where and when-influences on cutaneous vitamin D synthesis. *Prog Biophys Mol Biol* 2006;92:17–25.
- 53 Webb AR, Kazantzidis A, Kift RC, *et al.* Meeting vitamin D requirements in white Caucasians at UK latitudes: providing a choice. *Nutrients* 2018;10:497–3.
- 54 Webb AR, Kazantzidis A, Kift RC, *et al.* Colour counts: sunlight and skin type as drivers of vitamin D deficiency at UK latitudes. *Nutrients* 2018;10:457–3.
- 55 World Health Organisation. Food and nutrition tips during Self-Quarantine, 2020. Available: <http://www.euro.who.int/en/health-topics/health-emergencies/coronavirus-covid-19/novel-coronavirus-2019-ncov-technical-guidance/food-and-nutrition-tips-during-self-quarantine>
- 56 Uk Government. Nutrition Update. Coronavirus/COVID-19 Special Edition. (Number 16: April - June 2020), 2020. Available: [https://modgovuk.sharepoint.com/sites/defnet/JFC/Pages/Defence-nutrition-advisory-service-\(DNAS\)-.aspx](https://modgovuk.sharepoint.com/sites/defnet/JFC/Pages/Defence-nutrition-advisory-service-(DNAS)-.aspx)
- 57 Royal Osteoporosis Society. Vitamin D and bone health: a practical clinical guideline for patient management, 2019. Available: <https://www.guidelines.co.uk/musculoskeletal-and-joints/-ros-vitamin-d-and-bone-health-guideline/454558.article>
- 58 Bouillon R, Marcocci C, Carmeliet G, *et al.* Skeletal and extraskeletal actions of vitamin D: current evidence and outstanding questions. *Endocr Rev* 2019;40:1109–51.
- 59 UK Government. Advice on vitamin D. Available: <https://www.nhs.uk/conditions/vitamins-and-minerals/vitamin-d/>