

Empirical study of the 30-s chair-stand test as an indicator for musculoskeletal disorder risk of sedentary behaviour in Japanese office workers: a cross-sectional empirical study

Azusa Arimoto ¹, Shoko Ishikawa,² Etsuko Tadaka ¹

To cite: Arimoto A, Ishikawa S, Tadaka E. Empirical study of the 30-s chair-stand test as an indicator for musculoskeletal disorder risk of sedentary behaviour in Japanese office workers: a cross-sectional empirical study. *BMJ Nutrition, Prevention & Health* 2021;4:e000211. doi:10.1136/bmjnp-2020-000211

¹Department of Community Health Nursing, Yokohama City University, Yokohama, Kanagawa, Japan

²Kanazawa Welfare and Health Center, Yokohama City Office, Yokohama, Kanagawa, Japan

Correspondence to

Dr Etsuko Tadaka, Community Health Nursing, Yokohama City University, Yokohama, Kanagawa 236-0027, Japan; e_tadaka@yokohama-cu.ac.jp

Received 27 November 2020

Revised 2 March 2021

Accepted 3 March 2021

Published Online First

22 March 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

ABSTRACT

Objectives Sedentary behaviour among office workers and the risk of adverse health outcomes are public health problems. However, risk indicators for these outcomes require invasive biochemical examination. A proactive screening tool using a non-invasive, easy-to-use method is required to assess the risk focused on musculoskeletal health for primary prevention. However, middle-aged adults have insufficient awareness of musculoskeletal disorders. This study examined to determine whether the 30-s chair-stand test (CS-30) can be used as a proactive screening index for musculoskeletal disorder risk of sedentary behaviour in office workers.

Design Cross-sectional study using self-administered questionnaires and physical measurements.

Setting Four workplaces located in a metropolitan area of Japan.

Participants 431 Japanese office workers aged 20–64 years. 406 valid sets of results remained (valid response rate: 94.2%).

Primary and secondary outcome measures Musculoskeletal function was measured using the CS-30, quadriceps muscle strength. Receiver operating characteristic curve analysis was used to determine the sensitivity, specificity and optimal cut-off value for the CS-30. The risk of future incidence of musculoskeletal disorders was calculated using current quadriceps muscle strength.

Results In total participants, 47.0% were male and the mean sitting time in work duration was 455.6 min/day (SD=111.2 min). The mean lower limb quadriceps muscle strength was 444.8 N (SD=131.3 N). For the optimum cut-off value of 23 on the CS-30 for all participants, sensitivity was 0.809 and specificity was 0.231. For men, the optimum cut-off was 25, with a sensitivity of 0.855 and a specificity 0.172. For women, the optimum cut-off was 21, with a sensitivity of 0.854 and a specificity 0.275.

Conclusions Sensitivity was high, but specificity was insufficient. The CS-30 may be a potential proactive screening index for musculoskeletal disorder risk of sedentary behaviour, in combination with other indicators.

What this paper adds

- ▶ This empirical study examined the 30-s chair-stand test is a potential proactive screening index for musculoskeletal disorder risk, in combination with the assessment of other adverse health outcome indicators of sedentary behaviour.
- ▶ This study used objectively assessed musculoskeletal function using physical function test and the device.
- ▶ This study used a cross-sectional design, and could not clarify the relationship between quadriceps muscle strength change and the onset of lower limb musculoskeletal disorders.

BACKGROUND

Sedentary behaviour and a physical inactivity are major problems that need to be addressed, especially among adult office workers.^{1–4} Previous studies have reported that sedentary behaviour is related to negative health outcomes such as cardiovascular disease,^{5–8} diabetes,^{5 6 8} metabolic syndrome⁹ and musculoskeletal disorders.¹⁰ Musculoskeletal health is the foundation of a disability-free life. However, middle-aged adults have insufficient awareness of musculoskeletal disorders¹¹ and sedentary behaviour.

Sedentary behaviour is defined as any waking behaviour characterised by an energy expenditure of ≤ 1.5 metabolic equivalents while in a sitting or reclining posture.¹² Previous studies reported the sitting time of Japanese adults to be about 8 hours/day^{7 13}—the longest among 20 examined countries.¹³ Thus, middle-aged workers are a crucial population for assessing musculoskeletal function index as an indicator of sedentary behaviour.

Limited studies have identified the relationships between lower limb musculoskeletal disorders and sedentary behaviours

among office workers.^{14–17} Previous studies which were conducted in Japan revealed that 22%–40% of residents in their 40s and 25%–49% of those in their 50s had lower limb musculoskeletal disorders.^{18–19} These findings have highlighted that early detection and intervention for lower limb musculoskeletal disorders is required to sustain musculoskeletal function.

Musculoskeletal functions related to the lower limb musculoskeletal disorders¹⁸ are muscle strength¹⁴ and speed of motion.²⁰ These lower limb musculoskeletal functions among office workers have been assessed by using surface electromyography,^{16–21} muscle strength measurement using an optical measurement system¹⁷ and with the questionnaire, which includes subjective questions about pain, walking, lifestyle, personal care actions and social activities.¹¹

However, devices such as electromyography and optical measurement systems can produce objective assessments in experimental, non-daily situations but require a great deal of time. A questionnaire can be administered with little time and is easy to use; however, this assessment approach is subjective. An inexpensive and easy-to-use screening index is therefore needed to identify those who could benefit from primary prevention of musculoskeletal function decline among sedentary adult workers.

The 30-s chair-stand test (CS-30) has high test–retest reliability and high criterion-related validity with leg extension muscle strength.^{22–23} CS-30 was demonstrated by the test's ability to detect differences between various age and physical activity (PA)-level groups and CS-30 performance was significantly lower for low-active participants than for high-active participants.²² The CS-30 may therefore be usable as a new proactive screening index for musculoskeletal function decline as a measure of sedentary behaviour and may contribute to primary prevention in middle-aged individuals.

The objective of this study was to establish whether the CS-30 is a suitable proactive screening index for evaluating musculoskeletal disorder risk of sedentary behaviour in adult office workers.

METHODS

Study design and participants

The study used a cross-sectional design with anonymous, self-report questionnaires and physical measurements. The setting was two local government offices and two companies in a metropolitan area in Japan. The inclusion criteria for the participants were adult office workers aged 20–64 years who answered the questionnaires and provided physical measurements. Study participants were openly recruited via a flyer and poster at each worksite. Researchers and trained research public health nurses visited each worksite, administered a survey and physical measurements from September to November 2018.

Measurements

Questionnaires

Demographic characteristics

Demographic characteristics included age, gender, household composition, educational background and employment status.

Physical and mental characteristics

Participants responded 'yes' or 'no' to indicate whether they were currently being treated for each listed disease including musculoskeletal disorders. Participants with musculoskeletal disorders were excluded. Mental characteristics were measured using the Japanese version of the Kessler 6 (K6),²⁴ which consists of six items. Each item is scored on a 5-point Likert scale ranging from 0 to 4. The total possible score ranges from 0 to 24, with higher scores indicating higher depression and a K6 score above 5 identifies people at risk of depression.²⁴ Cronbach's alpha was 0.881 in this study.

Lifestyle characteristics

PA was measured using the Japanese version of the International Physical Activity Questionnaire-Short Form (IPAQ-SF),²⁵ a 7-day self-administered questionnaire.²⁶ The IPAQ-SF asks about three types of PAs: walking, moderate PA and vigorous PA. Computation of the total score requires the summation of the products of the duration (in minutes) and the frequency (in days) of walking, moderate PA and vigorous PA.²⁷ The IPAQ-SF is considered reliable and valid in 12 countries including Japan.²⁶

Commuting and employment situation characteristics that indicate a sitting time were measured using the Worker's Living Activity-time Questionnaire (WLAQ),^{4,28} which was designed by the National Institute of Occupational Safety and Health Japan (JNIOSH). The JNIOSH-WLAQ asked the participants about time spent sitting, standing and walking during their working and commuting time, typical domains of workers' lives.

Physical measurements

Body mass index

Body mass index (BMI) was calculated using height and body weight measurements. Height was measured with a stadiometer (seca 213; seca, Chiba, Japan) and body weight was measured with a digital weight scale (UC-322; A&D, Tokyo, Japan). Overweight was defined as $BMI \geq 25$, standard weight was defined as $18.5 \leq BMI < 25$ and low weight was defined as $BMI < 18.5$, based on the criteria.²⁹

Musculoskeletal function

The CS-30 involves repeatedly standing up from a seated position in a chair over a period of 30 s, as fast as possible.^{22–23} This test was developed by Jones *et al*²² to evaluate lower limb muscle strength. The test has high reliability using the test–retest method ($r=0.92$), as well as a strong correlation with leg extension strength (leg press; $r=0.71$).^{22–23} Nakatani *et al* modified the CS-30 and showed that this modified CS-30 can evaluate muscle strength for use in Japanese populations of all ages.^{23–30–31}

The participants were instructed to place one foot slightly in front of the other to help maintain balance when standing and to cross their arms at the wrists and hold them against the chest. The score was the total number of stands executed correctly within 30 s. Incorrectly executed stands were not counted.²²

Quadriceps muscle strength (QMS) was measured using the Locomo Scan (ALCARE, Tokyo, Japan). This device was developed based on the method for quadriceps setting training in a knee extension position.³² The Locomo Scan allows chronological measurement of knee extension muscle strength using strain gauge measurement. The smallest measurement unit was 1 N, and the maximum measured value was 1500 N.³² Omori *et al*³³ have previously reported that the maximum measured values using a prototype of Locomo Scan correlated significantly with the maximum measured values taken using an isokinetic machine (Biodex System 3; Biodex Medical System, Shirley, New York, USA).^{34 35} We calculated the weight bearing index (WBI) as QMS (in kilogram-force) divided by body weight (in kilograms).

Data analysis

IBM SPSS Statistics V.22.0 was used for the analysis. A critical endpoint calculated by the rates of change in QMS values affected the onset of lower limb musculoskeletal disorders.³⁶ Based on the reference QMS standardised values for adults,³² the rate of change in QMS between each 10-year age group from participants in their 20s to those in their 60s and reference values in older ages showing the onset of lower limb musculoskeletal disorders were calculated separately for men and women. The average QMS value in those showing musculoskeletal disorder onset was clarified by Takagi *et al*³⁶ and was used in this study as the QMS value related to the onset of musculoskeletal disorders as the critical endpoint. Those with QMS values below the values recommended for adults to avoid musculoskeletal disorders were categorised as showing musculoskeletal function decline indicating the adverse health outcome of sedentary behaviour (the risk group), and other participants were defined as the non-risk group. Among men, a score of 433 N or below in their 20s, a score of 430 N or below in their 30s, a score of 440 N or below in their 40s, a score of 359 N or below in their 50s and a score of 333 N or below in their ages 60–64 were classified as the risk group. Among women, a score of 419 N or below in their 20s, a score of 368 N or below in their 30s, a score of 334 N or below in their 40s, a score of 293 N or below in their 50s and a score of 242 N or below in their ages 60–64 were classified as the risk group. Receiver operating characteristic (ROC) curve analysis was used to compare the risk and CS-30 score to determine the sensitivity, specificity and validity of the optimal CS-30 cut-off value as a sedentary behaviour index for all participants, and for men and women separately, with cut-off values with 80% sensitivity based on Youden's Index.³⁷ Two-sided statistical tests were considered significant with an alpha level of 0.05.

Patient and public involvement

Patients or the public were not involved in the design or planning of the study.

RESULTS

A total of 431 people in the four workplaces agreed to participate. After the exclusion of 25 participants (19 participants because of incomplete information and 6 participants because of musculoskeletal disorders), 406 valid sets of results remained (valid response rate: 94.2%). The demographic characteristics of the participants are shown in [table 1](#). Their average age was 44.1 years (SD=10.2 years), 47.0% were male and 72.4% were full-time workers.

The mental and lifestyle characteristics of the participants are shown in [table 2](#).

In total, 68.5% had a low level of PA according to the IPAQ-SF. The mean one-way commute duration was 55.2 min (SD=26.6 min), and the mean work duration was 577.0 min/day (SD=83.6 min) according to the JNIOASH-WLAQ.

The mean sitting time in one-way commute duration was 20.1 min (SD=21.5 min), and the mean sitting time in work duration was 455.6 min/day (SD=111.2 min).

The participants' physical measurements are shown by sex in [table 3](#). The mean CS-30 score was 20.1 (SD=4.5), the mean lower limb QMS (either right or left) was 444.8 N (SD=131.3 N), the mean lower limb WBI (either right or left) was 0.74 (SD=0.22). The CS-30 was significantly weakly or moderately correlated with lower limb QMS ($r=0.293$ in total, 0.186 in men and 0.361 in women, $p<0.001$) and lower limb WBI ($r=0.305$ in total, 0.219 in men and 0.487 in women, $p<0.001$).

In total, 63 men (33.0%) were identified as having a high risk of declining musculoskeletal function. Of these, five participants in their 20s (21.7%) had a low QMS score of 433 N or below, 16 in their 30s (34.0%) had a QMS score of 430 N or below, 27 in their 40s (47.4%) had a QMS score of 440 N or below, 12 in their 50s (24.0%) had a QMS score of 359 N or below and 3 aged 60–64 years (21.4%) had a QMS score of 333 N or below. A total of 48 women (22.6%) had a high risk of declining musculoskeletal function. Of these, 7 in their 20s (33.3%) had a low QMS score of 419 N or below, 16 in their 30s (43.2%) had a QMS score of 368 N or below, 16 in their 40s (16.8%) had a QMS score of 334 N or below, 8 in their 50s (14.5%) had a QMS score of 293 N or below and 1 aged 60–64 years (14.2%) had a QMS score of 242 N or below. The ROC curves for the ability of the CS-30 to assess risk of declining musculoskeletal function as an indicator of sedentary behaviour are shown in [figures 1–3](#).

We determined the sensitivity, specificity and validity of the optimal CS-30 cut-off value for all participants, and for men and women separately, with cut-off values with 80% sensitivity based on Youden's Index.³⁷ Three cut-off values are shown for the ROC curve for each gender in addition to all participants because both the reference

Table 1 Participants' demographic characteristics

Items	All participants (n=406)		Male (n=191)		Female (n=215)		P value
	n or Mean±SD	% (Range)	n or Mean±SD	% (Range)	n or Mean±SD	% (Range)	
Age (years), n=406	44.1±10.2	(20–64)	43.8±11.0	(20–64)	44.3±9.3	(20–64)	0.598
20–29	44	10.8	23	11.9	21	9.8	0.018
30–39	84	20.8	47	24.6	37	17.3	
40–49	152	37.4	57	29.8	95	44.0	
50–59	105	26.0	50	26.3	55	25.5	
60–64	21	5.1	14	7.3	7	3.3	
Gender, n=406							
Male	191	47.0					
Female	215	53.0					
Household composition, n=404							0.724
Living alone	75	18.5	35	18.3	40	18.6	
Living with spouse	76	18.7	33	17.3	43	20.0	
Living with spouse and children	171	42.1	87	45.5	84	39.1	
Living with parents and children	25	6.2	11	5.8	14	6.5	
Others	57	14.0	24	12.6	33	15.3	
Educational background, n=404							0.000
Junior high school	2	0.5	1	0.5	1	0.5	
High school	38	9.4	20	10.5	18	8.4	
Junior college	65	16.0	9	4.7	56	26.0	
University	255	62.8	128	67.0	127	59.1	
Graduate school	43	10.6	31	16.2	12	5.6	
Other	1	0.2	1	0.5	0	0.0	
Employment status, n=389							0.000
Full time	294	72.4	168	88.0	126	58.6	
Part time	20	4.9	13	6.8	7	3.3	
Other	75	18.5	4	2.1	71	33.0	

QMS standardised values for adults³² and the average QMS values in those showing musculoskeletal disorder onset clarified by Takagi *et al*³⁶ differed by gender. For all participants, the optimum CS-30 cut-off was standing up 23 times, which had a sensitivity of 0.809 and a specificity of 0.231 with an area under the curve (AUC) of 0.565. For men, the optimum cut-off was standing up 25 times, with a sensitivity of 0.855 and a specificity of 0.172 with an AUC of 0.528. For women, the optimum cut-off was standing up 21 times, with a sensitivity of 0.854 and a specificity of 0.275 with an AUC of 0.637.

DISCUSSION

The physical measurement results for the lower limb revealed a mean QMS score of 444.8 N and a mean WBI score of 0.74. One previous study using the same device showed that the mean QMS for participants aged

from their 20s to their 60s was 485.7 N.³² Another study reported that a WBI score ranging from 0.6 to 0.8 in the general population indicates no hindrance in daily performance.³³ The present study's total sitting time per day is similar to a previous report of sedentary time among Japanese adult workers (mean: 8.4±3.4 hours/day)⁷ and longer than a previous study results for sitting time among Japanese adults (median: 420 min).³⁸ Therefore, participants in the present study may be similar to middle-aged workers in the general population in Japan. In both of the previous studies^{32,33} used to calculate the cut-off value, the middle-aged and older age groups included working people and non-working people. In a previous study¹⁹ that included community residents who were not working, 49% of participants in their 50s had lower limb musculoskeletal disorders. It has been shown that middle-aged and older workers have higher levels

Table 2 Participants' mental and lifestyle characteristics

Item	All participants (n=406)		Male (n=191)		Female (n=215)		P value
	n or Mean±SD	% (Range)	n or Mean±SD	% (Range)	n or Mean±SD	% (Range)	
Mental characteristics							
K6, n=404	3.0±3.7	(0–21)	3.0±3.8	(0–18)	2.9±3.6	(0–21)	0.696
Depression, K6 ≥5	111	27.6	49	43.8	62	29.3	0.473
Lifestyle characteristics							
Physical activity (IPAQ-SF), n=404							
Physical activity level							
Low	278	68.5	123	65.4	155	73.8	0.182
Moderate	108	27.1	58	30.9	50	23.8	
High	12	3.0	7	3.7	5	2.4	
Commuting and employment situation (JNIOOSH-WLAQ; min), n=405							
One-way commuting time	55.2±26.6	(5.0–155.0)	61.3±28.3	(10.0–155.0)	49.7±23.9	(5.0–120.0)	<0.001
Walking time	19.3±11.1	(0–90.0)	19.7±10.8	(0–60.0)	19.0±11.3	(0–90.0)	0.498
Sitting time	20.1±21.5	(0–110.0)	21.4±24.0	(0–110.0)	18.9±18.9	(0–110.0)	0.259
Standing time	13.9±16.8	(0–100.0)	17.3±19.1	(0–100.0)	10.8±13.7	(0–90.0)	<0.001
Working time	577.0±83.6	(265.0–815.0)	481.2±100.8	(300.0–815.0)	553.8±85.7	(110.0–800.0)	<0.001
Sitting time	455.6±111.2	(20.0–782.0)	481.2±100.8	(50.0–732.6)	433.0±115.2	(19.5–782.1)	<0.001
Standing time	120.7±86.8	(0–476.0)	122.0±82.2	(0–476.0)	118.5±90.6	(0–464.0)	0.774

IPAQ-SF, International Physical Activity Questionnaire-Short Form; JNIOOSH-WLAQ, National Institute of Occupational Safety and Health Japan-Worker's Living Activity-time Questionnaire; K6, Kessler 6.

Table 3 Participants' physical measurement results

Item	All participants (n=406)		Men (n=191)		Women (n=215)		P value
	n or Mean±SD	% (Range)	n or Mean±SD	% (Range)	n or Mean±SD	% (Range)	
Musculoskeletal function							
CS-30 (times)	20.1±4.5	(11–34)	20.9±4.4	(11–33)	19.3±4.4	(10.0–34.0)	<0.001
QMS (N)	444.8±131.3	(103.0–883.0)	471.6±146.1	(137.0–883.0)	421.0±111.6	(103.0–678.0)	<0.001
WBI	0.74±0.22	(0.00–2.00)	0.69±0.23	(0.23–1.56)	0.78±0.21	(0.18–1.34)	<0.001
Height (cm)	164.6±8.0	(148.0–183.0)	171.0±5.3	(153.2–183.1)	158.9±5.0	(146.9–173.2)	<0.001
Weight (kg)	62.4±12.8	(41.0–155.0)	70.4±12.2	(50.4–155.0)	55.3±8.6	(40.6–84.9)	<0.001
BMI (kg/m ²)	22.9±3.6	(15.2–46.5)	24.0±3.6	(18.0–46.5)	21.9±3.3	(15.2–35.7)	<0.001
Low weight, BMI<18.5	28	6.9	5	2.6	23	10.7	0.001
Standard weight, 18.5≤BMI<25	291	71.7	135	70.7	156	72.6	
Overweight, BMI≥25	87	21.4	51	26.7	36	16.7	

BMI, body mass index; CS-30, 30-s chair-stand test; QMS, quadriceps muscle strength; WBI, weight bearing index.




Figure 1 Receiver operating characteristic curves of 30-s chair-stand test (CS-30) as an indicator of sedentary lifestyle in all participants. CS-30 optimal cut-off: ≤ 23 . Sensitivity: 0.809. Specificity: 0.231. Area under the curve: 0.565.




Figure 3 Receiver operating characteristic curves of 30-s chair-stand test (CS-30) as an indicator of sedentary lifestyle in women. CS-30 optimal cut-off: ≤ 21 . Sensitivity: 0.854. Specificity: 0.275. Area under the curve: 0.637.

of health than non-workers. This suggests that workers above 50 years of age in the present study had better QMS values than the reference values.

A physical function index as an indicator of sedentary behaviour risk was estimated from the risk of future incidence of musculoskeletal disorders and optimum CS-30




Figure 2 Receiver operating characteristic curves of 30-s chair-stand test (CS-30) as an indicator of sedentary lifestyle in men. CS-30 optimal cut-off: ≤ 25 . Sensitivity: 0.855. Specificity: 0.172. Area under the curve: 0.528.

cut-off values with a sensitivity of over 0.80 were identified as 23 for all participants, 25 for men and 21 for women. These results suggest that there is a low probability of a false negative (determining no musculoskeletal disorder risk when the risk is actually high). The CS-30 could therefore be useful as a proactive screening test for identifying and targeting primary prevention programmes in the large population of healthy individuals with low levels of risk, as an effective public health strategy.³⁹ The specificity of these cut-off points was low, at about 0.20, which is similar to the findings of a previous study that reported a sensitivity of 1.000 when the specificity was 0.196 for using the seated toe-touch test to identify musculoskeletal disorders.³⁴ A performance test to be used before is more expensive and precise diagnostic test, the cut-off value had high sensitivity (0.900), whereas specificity was 0.260.⁴⁰ These findings suggest that the method of setting cut-off points with high sensitivity is valid. However, the specificity could have been low because we used a hard endpoint that only assessed risk of declining musculoskeletal function as an indicator of sedentary behaviour based on the risk of onset of lower limb musculoskeletal disorders and the QMS score.

The CS-30 only involves counting the number of times the person stands up in 30 s and no complicated calculations are necessary. This measurement is therefore easy to conduct and provides a realistic assessment of an individual's performance. This study used an examiner to ensure accuracy. However, in practice, self-checks using the second hand on a watch or a timer are also possible. The explanation and practice before the measurement are not time consuming because standing up is a normal

part of everyday behaviour. The CS-30 is also very safe and has been used successfully with frail older adults.^{22 23} The CS-30 is therefore a performance test that can be used to evaluate musculoskeletal function, as an indicator of sedentary behaviour. This method is simple and non-invasive. Even if an individual is judged to be at high risk of declining musculoskeletal function, any physical, mental or economic burden will be caused by the examination, diagnosis or treatment.

The present study had several limitations. First, this study focused on lower limb musculoskeletal disorder onset as an indicator of musculoskeletal function deterioration among the various negative outcomes caused by sedentary behaviour. Other indexes focusing on the additional outcomes of musculoskeletal functional decline can be selected as proactive screening indicators to understand the preliminary stages of musculoskeletal disease onset. Second, this study was cross-sectional in design. A longitudinal study is required to clarify the relationship between QMS change and the onset of lower limb musculoskeletal disorders, as well as to clarify the optimal predictive power of the CS-30 cut-off values identified in this study. Selection bias in this study may have occurred because the target population was limited to employees who worked for four Japanese companies and voluntarily responded to the open recruitment. For example, it is possible that people with a strong interest in health were more likely to cooperate, whereas those with lower limb muscle strength issues may have been less likely to take part. Future studies should be conducted in a wider range of countries, regions and occupations, and with a larger number of subjects. For the CS-30, we considered that the possibility of interviewer bias is relatively low due to training and unified procedures. Future studies will be needed to verify the inter-rater reliability of this method.

CONCLUSION

This study aimed to establish whether the CS-30 is a suitable proactive screening index for evaluating musculoskeletal disorder risk of sedentary behaviour in adult office workers. The results indicated that the optimal CS-30 cut-off value was 23 overall, 25 for men and 21 for women. Sensitivity was high, but specificity was insufficient. The CS-30 may be a potential proactive screening index for musculoskeletal disorder risk of sedentary behaviour, in combination with other indicators of other adverse health outcomes of sedentary behaviour to facilitate early primary prevention of adverse musculoskeletal health outcomes.

Acknowledgements The authors thank all the participants in this study. The authors also thank Assistant Professor K Shiratani, E Ito and all members of the Department of Community Health Nursing, Graduate School of Medicine, Yokohama City University. The authors thank Edanz Group (<https://en-author-services.edanz.com/ac>) for editing a draft of this manuscript.

Contributors AA, SI and ET developed the concept and design of this study. SI and ET were responsible for funding. AA and SI were responsible for the data collection, analysis, drafting and revising the manuscript. ET was responsible for obtaining

institutional review board approval, the supervision and for the reporting of study results. All authors read and approved the final manuscript.

Funding This work was supported by research grants from the Health Science Centre Foundation, Japan (Principal Investigator: SI).

Competing interests None declared.

Patient and public involvement statement Patients or public were not involved in the design or planning of the study.

Patient consent for publication Not required.

Ethics approval The Human Genome/Genetic Research Ethics Committee of the Medical Department of Yokohama City University, Japan (A180700001). The questionnaire in this study was unsigned to maintain the anonymity of all participant information. The researchers explained the contents of the survey to the participants, and consent was considered as consent to the survey by answering and submitting a questionnaire.

Provenance and peer review Not commissioned; externally peer reviewed by Dr Emmanuel Baah, University of North Carolina System.

Data availability statement Data may be obtained from a third party and are not publicly available. Requests for access to data should be addressed to the corresponding author.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Azusa Arimoto <http://orcid.org/0000-0002-8950-674X>

Etsuko Tadaka <http://orcid.org/0000-0002-4001-5018>

REFERENCES

- Bernaards CM, Hildebrandt VH, Hendriksen IJM. Correlates of sedentary time in different age groups: results from a large cross sectional Dutch survey. *BMC Public Health* 2016;16:1121.
- Saidj M, Menai M, Charreire H, et al. Descriptive study of sedentary behaviours in 35,444 French working adults: cross-sectional findings from the ACTI-Cités study. *BMC Public Health* 2015;15:379.
- Kurita S, Shibata A, Ishii K, et al. Social-ecological correlates of accelerometer-measured occupational sitting among Japanese desk-based workers. *BMC Public Health* 2019;19:1489.
- Matsuo T, So R, Takahashi M. Workers' physical activity data contribute to estimating maximal oxygen consumption: a questionnaire study to concurrently assess workers' sedentary behavior and cardiorespiratory fitness. *BMC Public Health* 2020;20:22.
- Wilmot EG, Edwardson CL, Achana FA, et al. Sedentary time in adults and the association with diabetes, cardiovascular disease and death: systematic review and meta-analysis. *Diabetologia* 2012;55:2895–905.
- Biswas A, Oh PI, Faulkner GE, et al. Sedentary time and its association with risk for disease incidence, mortality, and hospitalization in adults: a systematic review and meta-analysis. *Ann Intern Med* 2015;162:123–32.
- Honda T, Chen S, Kishimoto H, et al. Identifying associations between sedentary time and cardio-metabolic risk factors in working adults using objective and subjective measures: a cross-sectional analysis. *BMC Public Health* 2014;14:1307.
- Ekelund U, Steene-Johannessen J, Brown WJ, et al. Does physical activity attenuate, or even eliminate, the detrimental association of sitting time with mortality? A harmonised meta-analysis of data from more than 1 million men and women. *Lancet* 2016;388:1302–10.
- Honda T, Chen S, Yonemoto K, et al. Sedentary bout durations and metabolic syndrome among working adults: a prospective cohort study. *BMC Public Health* 2016;16:888.
- Marshall S, Gyi D. Evidence of health risks from occupational sitting: where do we stand? *Am J Prev Med* 2010;39:389–91.
- Seichi A, Hoshino Y, Doi T, et al. Development of a screening tool for risk of locomotive syndrome in the elderly: the 25-question geriatric locomotive function scale. *J Orthop Sci* 2012;17:163–72.
- Tremblay MS, Aubert S, Barnes JD, et al. Sedentary Behavior Research Network (SBRN) - Terminology Consensus Project process and outcome. *Int J Behav Nutr Phys Act* 2017;14:75.

- 13 Bauman A, Ainsworth BE, Sallis JF, *et al.* The descriptive epidemiology of sitting. A 20-country comparison using the International physical activity questionnaire (IPAQ). *Am J Prev Med* 2011;41:228–35.
- 14 del Pozo-Cruz B, Gusi N, Adsuar JC, *et al.* Musculoskeletal fitness and health-related quality of life characteristics among sedentary office workers affected by sub-acute, non-specific low back pain: a cross-sectional study. *Physiotherapy* 2013;99:194–200.
- 15 da Costa BR, Vieira ER. Risk factors for work-related musculoskeletal disorders: a systematic review of recent longitudinal studies. *Am J Ind Med* 2010;53:285–323.
- 16 Pesola AJ, Laukkanen A, Haakana P, *et al.* Muscle inactivity and activity patterns after sedentary time--targeted randomized controlled trial. *Med Sci Sports Exerc* 2014;46:2122–31.
- 17 Genin PM, Dessenne P, Finaud J, *et al.* Effect of work-related sedentary time on overall health profile in active vs. inactive office workers. *Front Public Health* 2018;6:279.
- 18 Yoshimura N, Muraki S, Nakamura K, *et al.* Epidemiology of the locomotive syndrome: the research on osteoarthritis/osteoporosis against disability study 2005-2015. *Mod Rheumatol* 2017;27:1–7.
- 19 Ito E, Tadaka E, Shiratani K, *et al.* Population-Based study to explore the risk factors to locomotive syndrome among community-dwelling middle-aged people in an urban area. *J Jpn Acad Com Health Nurs* 2017;20:4–11.
- 20 Arai T, Obuchi S, Shiba Y, *et al.* The validity of an assessment of maximum angular velocity of knee extension (Ke) using a gyroscope. *Arch Gerontol Geriatr* 2012;54:e175–80.
- 21 Baker R, Coenen P, Howie E, *et al.* The short term musculoskeletal and cognitive effects of prolonged sitting during office computer work. *Int J Environ Res Public Health* 2018;15:1678.
- 22 Jones CJ, Rikli RE, Beam WC. A 30-S chair-stand test as a measure of lower body strength in community-residing older adults. *Res Q Exerc Sport* 1999;70:113–9.
- 23 Nakatani T, Nadamoto M, Mimura K-I, *et al.* Validation of a 30-sec chair-stand test for evaluating lower extremity muscle strength in Japanese elderly adults. *Japan Journal of Physical Education, Health and Sport Sciences* 2002;47:451–61.
- 24 Furukawa TA, Kawakami N, Saitoh M, *et al.* The performance of the Japanese version of the K6 and K10 in the world mental health survey Japan. *Int J Methods Psychiatr Res* 2008;17:152–8.
- 25 Murase N, Katsumura T, Ueda C, *et al.* Validity and reliability of Japanese version of international physical activity questionnaire. *Journal of Health and Welfare Statistics* 2002;49:1–9.
- 26 Craig CL, Marshall AL, Sjöström M, *et al.* International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* 2003;35:1381–95.
- 27 IPAQ Research Committee. Guidelines for data processing and analysis of the International physical activity questionnaire (IPAQ) – short and long forms.p1-15, 2005. Available: <http://www.ipaq.ki.se/> [Accessed 13 Mar 2020].
- 28 Matsuo T, So R, Sasai H, *et al.* [Evaluation of Worker's Living Activity-time Questionnaire (JNIOH-WLAQ) primarily to assess workers' sedentary behavior]. *Sangyo Eiseigaku Zasshi* 2017;59:219–28.
- 29 Examination Committee of Criteria for 'Obesity Disease' in Japan, Japan Society for the Study of Obesity. New criteria for 'obesity disease' in Japan. *Circ J* 2002;66:987–92.
- 30 Nakatani T, Kawata Y, Nadamoto M. Effectiveness of a 30-second chair stand test to easily evaluate the leg muscle strength of young people. *Journal of health, physical education and recreation* 2002;52:661–5. Japanese.
- 31 Nakatani T, Nadamoto M, Mimura K. Aging change and standard value of 30-second chair stand test result. *Japanese Journal of Clinical Sports Medicine* 2003;20:349–335. Japanese.
- 32 Narumi K, Funaki Y, Yoshimura N, *et al.* Quadriceps muscle strength reference value as index for functional deterioration of locomotive organs: data from 3617 men and women in Japan. *J Orthop Sci* 2017;22:765–70.
- 33 Omori G, Koga Y, Tanaka M, *et al.* Quadriceps muscle strength and its relationship to radiographic knee osteoarthritis in Japanese elderly. *J Orthop Sci* 2013;18:536–42.
- 34 Bohannon RW. Reference values for extremity muscle strength obtained by hand-held dynamometry from adults aged 20 to 79 years. *Arch Phys Med Rehabil* 1997;78:26–32.
- 35 Nakamura M, Hashizume H, Oka H, *et al.* Physical performance measures associated with locomotive syndrome in middle-aged and older Japanese women. *J Geriatr Phys Ther* 2015;38:202–7.
- 36 Takagi S, Omori G, Koga H, *et al.* Quadriceps muscle weakness is related to increased risk of radiographic knee oa but not its progression in both women and men: the Matsudai knee osteoarthritis survey. *Knee Surg Sports Traumatol Arthrosc* 2018;26:2607–14.
- 37 Youden WJ. Index for rating diagnostic tests. *Cancer* 1950;3:32–5.
- 38 Kurita S, Shibata A, Ishii K, *et al.* Patterns of objectively assessed sedentary time and physical activity among Japanese workers: a cross-sectional observational study. *BMJ Open* 2019;9:e021690.
- 39 Zulman DM, Vijan S, Omenn GS, *et al.* The relative merits of population-based and targeted prevention strategies. *Milbank Q* 2008;86:557–80.
- 40 El Ghoch M, Rossi AP, Calugi S, *et al.* Physical performance measures in screening for reduced lean body mass in adult females with obesity. *Nutr Metab Cardiovasc Dis* 2018;28:917–21.