


Evaluation of the safety and tolerability of exogenous ketosis induced by orally administered free beta-hydroxybutyrate in healthy adult subjects

Lisa Isabel Pimentel-Suarez,^{1,2} Adrian Soto-Mota ^{1,2}

To cite: Pimentel-Suarez LI, Soto-Mota A. Evaluation of the safety and tolerability of exogenous ketosis induced by orally administered free beta-hydroxybutyrate in healthy adult subjects. *BMJ Nutrition, Prevention & Health* 2023;6:e000672. doi:10.1136/bmjnph-2023-000672

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjnph-2023-000672>).

¹Metabolic Diseases Research Unit, National Institute of Medical Sciences and Nutrition Salvador Zubiran, Mexico, Mexico

²School of Medicine, Tecnológico de Monterrey - Campus Ciudad de Mexico, Ciudad de Mexico, Mexico

Correspondence to

Dr Adrian Soto-Mota;
adrian.sotom@incmnsz.mx

Received 11 April 2023

Accepted 8 June 2023

Published Online First

6 July 2023



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

ABSTRACT

Beta-hydroxybutyrate (D-BHB) is a metabolite with intrinsic signalling activity that has gained attention as a potentially clinically useful supplement. There are available supplements for inducing ketosis: ketone salts, ketone esters and medium-chain triglycerides. Even when all of them raise D-BHB in the blood and all are safe and well tolerated, they significantly differ in their safety profile, their palatability and their price. A fourth and potentially interesting option is to use biologically identical D-BHB, which it is already commercially available in the USA (American Ketone) and Greater China (MedPHA). However, its safety and tolerability had not yet been documented in the scientific literature. We evaluated the safety and tolerability of orally administered free D-BHB in a gender and age-balanced sample of 24 asymptomatic and overtly healthy adults. No participant showed acid-base abnormalities or electrolyte abnormalities. Secondary symptoms were reported after only 6.2% of all drink takes and none of the reports described the symptom as 'severe'. The most frequently reported secondary effects (19/720 or 2.6%) were gastrointestinal discomfort, headache (7/720 or 1%) and loss of appetite (7/720 or 1%). No correlation between weight-adjusted dose and frequency of secondary symptoms was observed. Free D-BHB was a safe and well-tolerated intervention for inducing sustained exogenous ketosis. Being bioidentical, salt-free and lacking intermediate metabolites, this form of supplementation could have a larger safety spectrum than salt or alcohol-based exogenous ketones. More research is warranted to assess its clinical efficacy in those clinical scenarios in which achieving ketosis rapidly could be beneficial.

INTRODUCTION

Several ketogenic interventions have shown benefits in a wide variety of clinical scenarios.^{1–5} Ketosis can be induced in different ways: by 72 hours of fasting,⁶ with a carbohydrate-restricted⁷ diet, and within minutes using supplements.⁸ While endogenous ketosis interventions have proven useful in many illnesses such as diabetes,² psychiatric illnesses,⁹ migraine,¹⁰ and cognitive decline,¹¹ in some time-sensitive clinical scenarios where

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Ketogenic supplements could be useful therapeutic tools in certain time-sensitive circumstances. A previously unexplored but attractive option is using bioidentical beta-hydroxybutyrate (D-BHB).

WHAT THIS STUDY ADDS

⇒ Diluted free D-BHB is a safe and well-tolerated intervention for inducing sustained exogenous ketosis.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ Being bioidentical, salt and alcohol free, further research is warranted on the potential clinical uses of free-BHB.

ketosis could be beneficial (eg, heart failure¹² or sepsis¹³) endogenous ketosis is impractical as patients cannot wait 72 hours for achieving a physiologically relevant production.

Beta-hydroxybutyrate (D-BHB) is a metabolite with intrinsic signalling activity,¹⁴ therefore, it has gained attention as a potentially clinically useful supplement. There are three thoroughly studied ketone supplements for inducing ketosis: ketone salts,^{15–17} ketone esters¹⁸ and medium-chain triglycerides.¹⁹ Even when they all raise the D-BHB in the blood and all are safe and well tolerated, they significantly differ in their safety profile, palatability and price.^{8 20}

A fourth and potentially interesting option is to use biologically identical (D-BHB). Being salt-free and having no intermediate metabolites or precursors, it could have a broader clinical reach than previous ketone supplements (eg, patients with mineral load restrictions such as those living with heart failure¹² or chronic kidney disease^{21 22}). While it is already commercially available in the USA (American Ketone) and Greater China (MedPHA), its safety and tolerability had not yet been documented in the scientific




Figure 1 Study workflow.

literature. In this work, we evaluated the safety and tolerability of Free D-BHB as a ketogenic supplement in healthy adults.

MATERIALS AND METHODS

This study was preregistered in ClinicalTrials.gov on 22 October 2022, with the identifier NCT05584371. The study workflow can be observed in [figure 1](#).

A gender-balanced and age-balanced sample of asymptomatic adults (most of them, healthcare workers) without illnesses that caused acid-base imbalances were recruited and asked to ingest 10 grams of D-BHB, diluted in one litre of water and sweetened with a sugar substitute derived from the stevia plant already embedded in the preparation, every morning between 9:00 and 11:00 hours for 4 weeks. Women of childbearing age were recruited only if using a long-term method of contraception including any form of hormonal contraception and intrauterine devices. The exclusion criteria included adults on a ketogenic diet, carbohydrate-restricted diet or intermittent fasting.

The exogenous ketones used in our study were provided by a joint donation from American Ketone and MedPHA. This bioidentical D-BHB is produced with the acid hydrolysis of Poly-3Hydroxy-Butyrate²³ and is available in 60 mL bottles with 10 g of free D-BHB each. Participants were instructed to dilute their content in 1 L of water before consuming it.

As previously used in other studies on ketone supplement studies,¹⁸ the frequency and severity of symptoms (mild, moderate, severe) were assessed every 2 weeks with a self-administered open questionnaire to avoid priming positive and negative symptom reports and the intervention adherence was corroborated with qualitative urinary ketone strips (calculated as the proportion of urinary ketone strips with detectable acetoacetate). Venous blood gases were analysed at the baseline, 2 and 4 weeks later to rule out acid-base disorders. All blood samples were handled and analysed at the central lab of the National Institute of Medical Sciences and Nutrition Salvador

Zubiran adhering to their internationally certified standard operating procedures.

Being haemodynamically stable, and consistent with previous studies, venous blood samples were preferred over arterial ones. Since acid-base analysis assumes arterial values, all acid-base variables were therefore ‘arterialised’ by adding 0.04 to the measured pH and subtracting 5 from the measured partial pressure of carbon dioxide (pCO₂).²⁴ Because of Mexico City’s altitude, the normal range for pCO₂ is different than for most places in the world (30.2±3.4 mm Hg).²⁵

Data were analysed using R V.4.0.3 with R-Studio 2022.07.2+576. Blood gas analysis was performed using CreateTableOne::CreateContTable using default parameters. Individual trajectory plots were produced using Microsoft Office Excel. The sample size was calculated for achieving a balanced age and sex sample based on previously reported safety and tolerability studies of ketone supplements.¹⁸ All results are reported as mean±SD and the statistical significance cut-off was p<0.05.

RESULTS

In total, 24 adults were recruited and all of them finished their participation. Their demographic characteristics by sex can be observed in [table 1](#).

No participant showed abnormalities in their vital signs or acid-base assessments in any of their visits ([figure 2](#)). Additionally, all acid-base parameters were non-significantly different across all time points ([table 2](#)). Additionally, no electrolyte abnormalities in Na, K, Cl or ionised Ca were observed (shown in the publicly available online supplemental dataset).

Regarding tolerability and palatability, all participants responded to the self-reported questionnaires at least once and had an adherence above 90%. Only 44 out of the 720 total takes (6.2%) of the drink reported secondary symptoms. Of which, none of the reports described the symptom as ‘severe’, (17/720 or 2.3%) were ‘moderate’ and the remaining (27/720 or 3.9%) were ‘mild’ ([table 3](#)).

The most reported secondary effects (19/720 or 2.6%) were gastrointestinal discomfort, headache (7/720 or 1%) and loss of appetite (7/720 or 1%). The rest of the reported secondary symptoms were reported <1% of the time. No participant reported more than two secondary symptoms nor the same symptom more than twice. No correlation between weight-adjusted dose and frequency of secondary symptoms was observed ([figure 3](#)).

DISCUSSION

Hereby and as observed with other formulations,^{16,18} exogenous ketosis with free D-BHB was safe and well tolerated. Since free-D-BHB has some advantages over other formulations, more research on its potential clinical applications is warranted.

Table 1 Baseline characteristics of the participants

	Women (n=13)	Men (n=11)
Age (years)	42.2±13.7	41.5±14.7
Height (m)	1.6±0.04	1.77±0.04
Weight (kg)	64.3±8.8	78.9±14.8
Dose (mg/kg)	158±20.9	129.8±19.2
Body mass index	23.9±3.2	20.9±6.3

Data are means±SD.

For example, ketone salts may not be suitable for patients who should restrict their salt intake, such as those with heart failure or kidney disease.²⁶ On the other hand, the ketone monoester has an alcohol precursor that could limit its use in patients with liver disease and has poor palatability.^{3,20} More importantly, being bioidentical, it is reasonable to assume that the safety and pharmacokinetic properties of free D-BHB are comparable to the endogenously produced metabolite.

It is worth reminding that since Mexico City is more than 2200 m above sea level, the arterial gasometric normal values are slightly different, and it is expected to observe lower CO₂ and HCO₃ levels in healthy individuals. This difference is relevant because clinicians elsewhere could incorrectly diagnose respiratory alkalosis with CO₂

Table 2 Acid-base analyses across all time points (arterialised values)

	Visit 1	Visit 2	Visit 3	P value
pH	7.41±0.03	7.43±0.03	7.44±0.03	0.19
pCO ₂	39.2±4.3	38.5±5.8	37±4.7	0.30
HCO ₃ ⁻	24.9±2.6	24.8±2.3	24.4±1.9	0.72

Data are means±SD. Hypothesis testing used a Kruskal-Wallis test.

values that are normal for people living in Mexico City.²⁵ Consistently with other ketone supplement formulations, even those inducing acute documented acid-base disturbances,⁸ the chronic ingestion of free D-BHB did not cause chronic acid-base or electrolyte alterations.^{8,15,16,18}

Among the strengths of this study, we should mention that most participants were healthcare workers and adherence was above 90%, making it unlikely that the low frequency of adverse effects is due to lack of adherence, and the close communication with participants. Moreover, since all participants consumed the same dose but have different weights a dose/kg spectrum is created allowing us to see if the frequency of adverse symptoms is more frequent in leaner individuals. As with other ketogenic supplements, no correlation was observed.¹⁸

Furthermore, we examined a sample size that is comparable to what has been examined for other commercially available ketogenic compounds. However, in contrast with those studies, we were able to enlist a more diverse age and gender distribution in our study. For example, Holland *et al*,¹⁵ who only studied individuals between the ages of 18 and 35, and Stefan *et al*¹⁷ focused exclusively on teenagers. This difference is particularly relevant as some of the potential clinical applications of exogenous ketones like heart failure¹² or cognitive decline¹¹ are on illnesses that are more frequent among elders.

Acid-base and blood electrolytes are the only markers that different types of exogenous ketones have shown to induce changes on.⁸ Therefore, when comparing studies evaluating the safety of different exogenous ketones, it is important to consider that some of the relevant markers, they measured are specific to the chemical nature of each supplement. In other words, due to their alcohol content, liver enzyme measurements are relevant for butanediol or



Figure 2 The individual trajectory of acid-base parameters across all time points. (A) pH, (B) pCO₂=partial pressure of carbon dioxide, (C) HCO₃=bicarbonate.

Table 3 Secondary symptoms following each of the 720 administered ketone drinks

Symptom	No of participants	Events* (mild/moderate)
Gastrointestinal discomfort	8	20/720 (10/10)
Loss of appetite	5	12/720 (9/3)
Headache	6	12/720 (8/4)

*Some participants recorded the same symptom more than once, so the occurrence was expressed relative to 720, the total number of drinks consumed (24×30 drinks).




Figure 3 Correlation between secondary symptom frequency and weight-adjusted dose.

the ketone ester but not for bioidentical BHB. Moreover, despite having used BHB doses many times greater than the one we worked with, none of the exogenous ketone supplements has been shown to induce detrimental changes in liver or kidney function tests.^{8 10 15 16 18} Notably, the only case where other biomarkers have been shown to change after consuming a ketone supplement, are migraine patients with baseline metabolic risk markers, in whom these markers (ie, C reactive protein, cortisol insulin and thyroid markers) improved.¹⁰

On the other hand, although comparable to what has been documented with other forms of ketone supplementation,^{8 10 15 16 18} we should mention its follow-up time among its limitations. Additionally, is not possible to make direct comparisons about palatability between the other available ketogenic supplements, and more studies directly addressing this question are required.²⁰

CONCLUSIONS

Diluted free D-BHB was a safe and well-tolerated intervention for inducing sustained exogenous ketosis. Being bioidentical, salt-free and lacking intermediate metabolites, this form of supplementation has a larger safety spectrum than salt or alcohol-based exogenous ketones. More research is warranted to assess its clinical efficacy in those clinical scenarios where achieving ketosis rapidly could be beneficial.

Acknowledgements The authors thank the Central Laboratory staff at the National Institute of Medical Sciences and Nutrition Salvador Zubiran for their invaluable support to make this work possible.

Contributors LIP-S recruited participants and collected data, and AS-M designed the study and analysed data. Both authors redacted this draft and approved the current version of the manuscript. AS-M is the guarantor of this work.

Funding The exogenous ketones used in our study were provided by a joint donation from American Ketone and MedPHA. However, the authors designed, carried out, analysed all data and wrote this report independently.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants and ethical approval was obtained from the Ethics Committee of the National Institute of Medical Sciences and Nutrition Salvador Zubirán with the file code UIE-4275-22-23-1 on 22 September 2022. Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available in a public, open access repository. <https://github.com/AdrianSotoM/FreeBHB>

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID ID

Adrian Soto-Mota <http://orcid.org/0000-0002-9173-7440>

REFERENCES

- Norwitz NG, Jaramillo JG, Clarke K, *et al*. Ketotherapeutics for neurodegenerative diseases. *Int Rev Neurobiol* 2020;155:141–68.
- Nicholas AP, Soto-Mota A, Lambert H, *et al*. Restricting carbohydrates and calories in the treatment of type 2 diabetes: a systematic review of the effectiveness of 'low-carbohydrate' interventions with differing energy levels. *J Nutr Sci* 2021;10:e76.
- Soto-Mota A, Norwitz NG, Evans RD, *et al*. Exogenous D-B-Hydroxybutyrate lowers blood glucose in part by decreasing the availability of L-alanine for gluconeogenesis. *Endocrinol Diabetes Metab* 2022;5:e00300.
- Norwitz NG, Hu MT, Clarke K. The mechanisms by which the ketone body D-B-Hydroxybutyrate may improve the multiple cellular pathologies of Parkinson's disease. *Front Nutr* 2019;6:63.
- Norwitz NG, Sethi S, Palmer CM. Ketogenic diet as a metabolic treatment for mental illness. *Curr Opin Endocrinol Diabetes Obes* 2020;27:269–74.
- Cahill GF, George FC. Fuel metabolism in starvation. *Annu Rev Nutr* 2006;26:1–22.
- Feinman RD, Pogozelski WK, Astrup A, *et al*. Dietary carbohydrate restriction as the first approach in diabetes management: critical review and evidence base. *Nutrition* 2015;31:1–13.
- Stubbs BJ, Cox PJ, Evans RD, *et al*. On the metabolism of exogenous ketones in humans. *Front Physiol* 2017;8:848.
- Danan A, Westman EC, Saslow LR, *et al*. The ketogenic diet for refractory mental illness: a retrospective analysis of 31 Inpatients. *Front Psychiatry* 2022;13:951376.
- Gross EC, Putananickal N, Orsini A-L, *et al*. Defining metabolic migraine with a distinct subgroup of patients with suboptimal inflammatory and metabolic markers. *Sci Rep* 2023;13:3787.
- Myette-Côté É, Soto-Mota A, Cunnane SC. Ketones - potential to achieve brain energy rescue and sustain cognitive health during aging. *Br J Nutr* 2022;128:407–23.
- Nielsen R, Møller N, Gormsen LC, *et al*. Cardiovascular effects of treatment with the ketone body 3-Hydroxybutyrate in chronic heart failure patients. *Circulation* 2019;139:2129–41.
- Soto-Mota A, Norwitz NG, Clarke K. Why a D-B-Hydroxybutyrate monoester? *Biochem Soc Trans* 2020;48:51–9.
- Rojas-Morales P, Tapia E, Pedraza-Chaverri J. B-Hydroxybutyrate: a signaling metabolite in starvation response? *Cell Signal* 2016;28:917–23.
- Holland AM, Qazi AS, Beasley KN, *et al*. Blood and cardiovascular health parameters after supplementing with ketone salts for six weeks. *Journal of Insulin Resistance* 2019;4:47.
- Buga A, Kackley ML, Crabtree CD, *et al*. The effects of a 6-week controlled, hypocaloric ketogenic diet, with and without exogenous ketone salts, on body composition responses. *Front Nutr* 2021;8:618520.
- Stefan M, Sharp M, Gheith R, *et al*. The effect of exogenous beta-hydroxybutyrate salt supplementation on metrics of safety and health in adolescents. *Nutrients* 2021;13:854.
- Soto-Mota A, Vansant H, Evans RD, *et al*. Safety and tolerability of sustained exogenous ketosis using ketone Monoester drinks for 28 days in healthy adults. *Regul Toxicol Pharmacol* 2019;109:104506.

- 19 St-Onge M-P, Jones PJH. Physiological effects of medium-chain triglycerides: potential of obesity. *The Journal of Nutrition* 2002;132:329–32.
- 20 Stubbs BJ, Cox PJ, Kirk T, *et al.* Gastrointestinal effects of exogenous ketone drinks are infrequent, mild and vary according to ketone compound and dose. *Int J Sport Nutr Exerc Metab* 2019;29:596–603.
- 21 Torres JA, Kruger SL, Broderick C, *et al.* Ketosis ameliorates renal cyst growth in polycystic kidney disease. *Cell Metab* 2019;30:1007–23.
- 22 Rojas-Morales P, Pedraza-Chaverri J, Tapia E. Ketone bodies for kidney injury and disease. *Advances in Redox Research* 2021;2:100009.
- 23 Seebach D, Beck A, Breitschuh R. Direct degradation of the biopolymer Poly[(R)-3-Hydroxybutyric acid] to (R)-3-Hydroxybutanoic acid and its methyl ester. *Org Synth* 1993;71:39.
- 24 Walkey AJ, Farber HW, O'Donnell C, *et al.* The accuracy of the central venous blood gas for acid-base monitoring. *J Intensive Care Med* 2010;25:104–10.
- 25 Cid-Juárez S, Téllez-Navarrete NA, Bautista-Bernal A, *et al.* Arterial blood gases in normal subjects at 2240 meters above sea level: impact of age, gender, and body mass index. *Rev Invest Clin* 2023;75:29–36.
- 26 Borlaug BA. Evaluation and management of heart failure with preserved ejection fraction. *Nat Rev Cardiol* 2020;17:559–73.