

Title: Covid-19 illness in relation to sleep and burnout

Hyunju Kim PhD, Sheila Hegde MD, Christine LaFiura BA, Madhunika Raghavan MA, Eric Luong MPH, Susan Cheng MD, Casey M Rebholz PhD, Sara B Seidelmann MD PhD

Supplemental Table 1. Adjusted Odds Ratios (OR) and 95% Confidence Intervals (CI) for the association between Sleep and Duration and Severity of Covid-19 like illness (n=568)^a

Supplemental Table 2. Adjusted Odds Ratios (OR) and 95% Confidence Intervals (CI) for the Association between Sleep and Covid-19 test (PCR or antibody)^a

Supplemental Table 1. Adjusted Odds Ratios (OR) and 95% Confidence Intervals (CI) for the association between Sleep and Duration and Severity of Covid-19 like illness (n=568)^a

	> 14 days of symptoms ^b			Moderate to severe symptoms ^c		
	OR (95% CI)	P-value	P-trend	OR (95% CI)	P-value	P-trend
Number of sleeping hours at night	0.92 (0.76, 1.12)	0.42	--	0.83 (0.70, 0.99)	0.03	--
Number of daytime napping hours	0.93 (0.82, 1.05)	0.25	--	1.06 (0.97, 1.17)	0.21	--
Sleep disturbance						
Difficulty sleeping at night (<i>yes</i>)	1.13 (0.68, 1.88)	0.63	--	1.04 (0.65, 1.67)	0.87	--
Often wake up at early hours and unable to get back to sleep (<i>yes</i>)	0.98 (0.60, 1.60)	0.93	--	0.86 (0.54, 1.37)	0.53	--
Sleeping pill use >3 times/wk (<i>yes</i>)	1.35 (0.59, 3.07)	0.48	--	1.28 (0.6, 2.72)	0.52	--
Self-reported sleep disturbance score						
No sleep problem						
Self-report of having 1 sleep problem	0.90 (0.49, 1.67)	0.74	--	0.59 (0.32, 1.11)	0.10	--
Self-report of having 2 sleep problems	1.21 (0.66, 2.22)	0.54	--	1.06 (0.6, 1.88)	0.84	--
Self-report of having 3 sleep problems	1.05 (0.37, 2.94)	0.93	--	1.09 (0.44, 2.71)	0.85	--
Any sleep disturbance (≥1 sleep problem)	1.04 (0.66, 1.65)	0.86	--	0.84 (0.54, 1.29)	0.43	--
Frequency of self-reported burnout			0.02			0.01
Never						
Rarely	0.93 (0.47, 1.82)	0.83	--	1.00 (0.55, 1.83)	0.99	--
Weekly	1.35 (0.59, 3.06)	0.47	--	1.47 (0.69, 3.12)	0.32	--
Every day	2.98 (1.10, 8.05)	0.03	--	3.26 (1.25, 8.48)	0.02	--

^a adjusted for age, sex, race, country, specialty, presence of a medical condition, stress from work (adjusted only for sleep variables), frequency of contact with a Covid-19 patient at work, close exposure to a Covid-19 patient inside the workplace without PPE and close exposure outside of the workplace

^b Participants reported the number of days they experienced symptoms of Covid-19. Asymptomatic individuals with a positive PCR or antibody test were considered to have 0 days of Covid-19 symptom duration.

^c Moderate to severe severity was compared to very mild to mild severity. 'Very mild' severity was defined as asymptomatic or nearly asymptomatic. 'Mild' severity was defined as symptoms [fever <38°C (without treatment), with or without cough, no dyspnea, no gasping, no abnormal imaging findings]. 'Moderate' severity was defined as fever, respiratory symptoms, and/or imaging findings of pneumonia. 'Severe'

severity was defined as meet any of the following: 1) respiratory distress, Respiratory Rate ≥ 30 times/min 2) low oxygen saturation (SpO₂) <93% at rest 3) partial pressure of oxygen (PaO₂)/ fraction of inspired oxygen (FiO₂) ≤ 300 mm Hg.

Supplemental Table 2. Adjusted Odds Ratios (OR) and 95% Confidence Intervals (CI) for the Association between Sleep and Covid-19 test (PCR or antibody)^a

	OR (95% CI)								
	Model 1 ^b	P-value	P-trend	Model 2 ^c	P-value	P-trend	Model 3 ^d	P-value	P-trend
Number of sleeping hours at night	0.89 (0.80, 1.01)	0.08	--	0.91 (0.81, 1.01)	0.09	--	0.91 (0.81, 1.02)	0.12	--
Number of daytime napping hours	1.00 (0.93, 1.07)	0.96	--	1.00 (0.93, 1.07)	0.99	--	1.00 (0.93, 1.07)	0.98	--
Sleep disturbance									
Difficulty sleeping at night (<i>yes</i>)	0.93 (0.67, 1.26)	0.63	--	0.88 (0.64, 1.21)	0.43	--	0.86 (0.62, 1.19)	0.36	--
Often wake up at early hours and unable to get back to sleep (<i>yes</i>)	0.85 (0.63, 1.15)	0.30	--	0.81 (0.59, 1.10)	0.17	--	0.77 (0.56, 1.05)	0.10	--
Sleeping pill use >3 times/week (<i>yes</i>)	1.25 (0.75, 2.12)	0.39	--	1.21 (0.72, 2.05)	0.47	--	1.15 (0.68, 1.95)	0.60	--
Self-reported sleep disturbance score									
No sleep problem	Ref	--	0.95	Ref	--	0.34	Ref	--	0.24
Self-report of having 1 sleep problem	0.81 (0.57, 1.14)	0.23	--	0.78 (0.55, 1.10)	0.17	--	0.75 (0.53, 1.07)	0.11	--
Self-report of having 2 sleep problems	0.70 (0.46, 1.07)	0.10	--	0.65 (0.42, 1.00)	0.05	--	0.63 (0.41, 0.97)	0.04	--
Self-report of having 3 sleep problems	1.77 (0.94, 3.33)	0.07	--	1.64 (0.87, 3.11)	0.12	--	1.50 (0.79, 2.85)	0.21	--
Any sleep disturbance (≥1 sleep problem)	0.84 (0.64, 1.09)	0.19	--	0.79 (0.60, 1.04)	0.11	--	0.76 (0.58, 1.01)	0.06	--
Frequency of self-reported burnout									
Never	Ref	--	0.02	Ref	--	0.02	Ref	--	0.04
Rarely	1.46 (1.00, 2.12)	0.05	--	1.46 (1.00, 2.13)	0.05	--	1.45 (0.99, 2.10)	0.05	--
Weekly	1.36 (0.85, 2.17)	0.19	--	1.36 (0.85, 2.17)	0.25	--	1.30 (0.82, 2.09)	0.32	--
Every day	2.80 (1.45, 5.39)	0.009	--	2.71 (1.46, 5.43)	0.002	--	2.60 (1.34, 5.05)	0.005	--

^a Covid-19 cases are defined as individuals with positive PCR or antibody test (N=298 cases and N=2316 controls).

^b Model 1 adjusted for age, sex, race, and country.

^c Model 2 additionally adjusted for specialty, presence of a medical condition, and stress from work (feeling burned out; only adjusted for sleep habits).

^d Model 3 additionally adjusted for frequency of contact with a Covid-19 patient at work, close exposure to a Covid-19 patient inside the workplace without PPE and close exposure outside of the workplace.

Bold font denotes statistically significant associations.