

Supplemental Table 1: The risk of incident hypertension within 10 years following delivery by servings per day of sucrose-sweetened beverages: MoBa Cohort Study (N= 60,027)

Entire cohort Servings of SSB ¹	Participants N	Cases n	Model 1 ² HR (95% CI)	Model 2 ³ HR (95% CI)	Model 3 ⁴ HR (95% CI)
<1	51,143	1,179	Referent	Referent	Referent
1-1.49	3728	122	1.40 (1.16, 1.69)	1.37 (1.13, 1.67)	1.34 (1.11, 1.64)
≥ 1.50	5156	179	1.55 (1.31, 1.82)	1.29 (1.09, 1.53)	1.23 (1.03, 1.48)

Abbreviations: CI, confidence interval; HR, hazard ratio; SSB, sugar-sweetened beverages.

1. One serving =250ml
2. Maternal age at delivery (years), total energy intake (kcal/day) evaluated in Cox proportional hazards analyses.
3. Maternal age at delivery (years), total energy intake (kcal/day), body mass index (kg/m²), daily smoking prior to pregnancy, parity, diabetes mellitus (prior or during pregnancy or type unspecified), gestational hypertension (with or without proteinuria), preterm delivery (<37 weeks gestation) and physical activity (≥3 hrs, <3 hrs per/week, missing).
4. Model 2 covariates plus intake of dietary fiber (g/day) and added sugar from food as percent energy.